

Ethics committee approval: This study was approved by the Ethics and Research Committee from Faculdade de Ceilândia (CEP/FCE), with CAAE number 80704617.5.0000.8093.

<https://doi.org/10.1016/j.bjpt.2024.100813>

217

CLINICAL AND SOCIODEMOGRAPHIC PROFILE OF WOMEN WHO HAD VAGINAL BIRTH IN A PUBLIC HOSPITAL IN DISTRITO FEDERAL: A PRELIMINARY STUDY

Julia Shimohara Bradaschia¹, Mariana Martinez David¹, Sthefany da Silva Souza¹, Serise Amaral Pequeno¹, Raquel Henriques Jácomo², Aline Teixeira Alves¹

¹ Universidade de Brasília (UnB), Faculdade de Ceilândia, Brasília, Distrito Federal, Brasil

² Hospital Universitário de Brasília, Universidade de Brasília, Brasília, Distrito Federal, Brasil

Background: In 2020, 47,44% of deliveries in Distrito Federal were vaginal according with the Live Birth Information System (SINASC). The cesarean sections exceeded three times the rate recommended by the WHO of 15%. To encourage the vaginal delivery in Brasil, programs, such as Rede Cegonha of SUS, were implemented to reinforce the importance of vaginal birth and the improvement in prenatal care until 2 years after birth. This study accompanies the characteristics and impacts of vaginal birth.

Objectives: Describe the vaginal birth parturient's profile from a public hospital in Distrito Federal from January to June of 2020.

Methods: It's an observational and cross-sectional study, that search through secondary data to describe the parturient's sociodemographic and clinical profile that gave birth between January and June of 2020. The inclusion criteria were vaginal birth with living newborn. Twin birth and illegible medical records were the exclusion criteria.

Results: Six hundred medical records met the inclusion criteria. The most prevalent group age for vaginal birth was 52,83% of women between 20 and 29 years old, with adolescents (14 to 19 years old) representing 12,83%. To identify the social demographic profile, about 65% identified themselves as brown-skinned, 52% were single and 45,67% were from Águas Lindas de Goiás. Regarding educational level, 52,33% had completed high school and 94,67% of the deliveries occurred with a companion. Concerning clinical data, 63% of the women were multiparous, 90,17% were over 37 weeks of pregnancy, 38% of deliveries were induced and 26,50% used some kind of anesthesia. As for childbirth positions, 68,67% were on the semi-sitting position, 92,67% were not instrumented and 56,67% had perineal lacerations being the second-degree – laceration of the vaginal mucosa and muscles of the pelvic floor being the most prevalent (32,50%). In only 3,67% episiotomy was performed. The average weight of the newborn was of 3,18 Kg, with 90,16% weighting more than 2500g. The average length, head circumference and first APGAR of newborns were 48,46 cm, 34,18 cm and 8,26, respectively. **Conclusion:** Most of the vaginal births were at term, and the most frequent birth position was semi-sitting. Regardless it's a Distrito Federal's public hospital, almost half the patients were from Goiás state. It is worth mentioning that adolescent between 14 to 19 years old maintains a percentage over 10% in the last 3 years of data in the first 6 months. The perineal injuries are still prevalent in more than 50% of the cases, and studies about the possible factors should be done.

Implications: With the presented data, it is noticed the need of implement actions about sex education and family planning to the teenager public, given that there's a significant percentile of parturient in this age group. Furthermore, it should be sought to

understand the high index of women that suffered perineal injuries in this period, to implement preventive measures.

Keywords: Pregnant Women, Natural Childbirth, Health Profile

Conflict of interest: The authors declare no conflict of interest.

Acknowledgment: The authors appreciate the Conselho Nacional de Desenvolvimento Científico e Tecnológico and the Fundação de Apoio à Pesquisa do Distrito Federal for the financial support.

Ethics committee approval: This study was approved by the Ethics and Research Committee from Faculdade de Ceilândia (CEP/FCE), with CAAE number 80704617.5.0000.8093.

<https://doi.org/10.1016/j.bjpt.2024.100814>

218

PROFILE OF HEALTHY HABITS OF INDIVIDUALS AFTER STROKE AND THEIR KNOWLEDGE ABOUT THIS HEALTH CONDITION: PRELIMINARY RESULTS

Júlia Stephanie Ruas dos Santos¹, Paula da Cruz Peniche¹, Jéssica Melo dos Santos¹, Maria Eduarda Matheus Lopes¹, Dâmaris Marçal Rodrigues¹, Christina Danielli Coelho de Moraes Faria¹

¹ Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, Minas Gerais, Brasil

Background: The recurrent stroke presents a high burden, mainly in developing countries, such as Brazil. Secondary prevention guidelines guide the adoption of a healthy lifestyle and having knowledge about stroke to control risk factors for recurrent stroke. Therefore, it is important to identify the profile of healthy habits and knowledge to direct the use of related interventions.

Objectives: To describe the profile of healthy habits and knowledge about stroke among individuals after stroke in a Brazilian metropolis.

Methods: Cross-sectional study, which recruited individuals two years after the stroke, aged ≥ 20 years, that received their first attendance in a stroke unit in the metropolis of Belo Horizonte (MG). For data collection, questions from the App "Stroke Riskometer" and from previous studies were used. Data were collected via telephone call and operationalized through absolute and relative frequency.

Results: Seventy-three individuals (63 ± 15 years old, 52% male) were included until now, and 7 (10%) cases of recurrent stroke were identified. As for habits, 64 (88%) reported not smoking, of these, 40 (55%) stopped smoking more than a year ago and 24 (33%) never smoked. Regarding alcohol consumption, 54 (74%) reported do not consume and, of these, 32 (59%) stopped consuming and 22 (41%) never consumed. As for eating, just over half ($n=48$, 66%) reported have adequate eating and 43 (59%) consume 2-3 fruits or vegetables/day. As for the physical activity practice, 29 (40%) reported practice it and, of these, 15 (52%) practice 2.5 hours/week and 11 (38%) between 1-2 hours/week. Just over half ($n=45$, 62%) reported to have not experienced significant emotional stress/depression for a year. As for knowledge, just over half reported know what a stroke is ($n=42$, 58%) and its signs and symptoms: difficulty speaking/understanding/articulating speech ($n=48$, 66%), loss of strength in arm ($n=48$, 66%) and smile/crooked mouth/face asymmetry ($n=56$, 77%). Most said they knew that smoking ($n=62$, 85%), alcohol consumption ($n=55$, 75%), inadequate eating ($n=57$, 78%), not practicing physical activity ($n=59$, 81%) and mental stress ($n=60$, 82%) are risk factors for stroke.

Conclusion: The habits that need more attention are healthy eating, alcohol consumption and mainly physical activity. Secondary prevention actions should also promote knowledge about what