



Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>



ORIGINAL RESEARCH

Does low and heavy load resistance training affect musculoskeletal pain in overweight and obese women? Secondary analysis of a randomized controlled trial[☆]



Anne Mette Rustaden^{a,*}, Lene Annette Hagen Haakstad^a, Gøran Paulsen^b, Kari Bø^a

^a Department of Sports Medicine, Norwegian School of Sports Sciences, Oslo, Norway

^b The Norwegian Olympic and Paralympic Committee and Confederation of Sport, Oslo, Norway

Received 3 November 2018; accepted 9 January 2019

Available online 18 January 2019

KEYWORDS

Strength training;
Group exercise;
Personal trainer;
Obesity;
Pain

Abstract

Background: Overweight and obesity are associated with musculoskeletal pain, particularly in the female population. However, regular resistance training may positively affect these complaints.

Objective: The present study aimed to investigate between group differences in musculoskeletal pain in previously inactive women, allocated to three different resistance-training modalities available in health- and fitness clubs.

Methods: This is secondary analysis from a single-blinded randomized controlled trial, including healthy women (aged 18–65) with a BMI (kg/m²) ≥ 25 . The participants were allocated to 12 weeks (3 times/weekly) of either BodyPump (high-repetition low-load group session) ($n = 24$), heavy load resistance training with a personal trainer ($n = 28$), non-supervised heavy load resistance training ($n = 19$) or non-exercising controls ($n = 21$). Primary outcome was self-reported musculoskeletal pain in ten different body parts, measured with the Standardized Nordic Pain Questionnaire, at baseline and post-test. In addition, the study included sub-analyses of the participants when they were divided into high (≥ 28 of 36 sessions, $n = 38$) and low (≤ 27 of 36 sessions, $n = 22$) exercise adherence.

Results: The analysis revealed no between group differences in musculoskeletal pain in any of the ten body parts. The results did not change when the participants were divided into high versus low adherence.

[☆] This paper is part of a Special Issue on Women's Health Physical Therapy.

* Corresponding author at: Department of Sports Medicine, Norwegian School of Sports Sciences, P.O. Box 4014, Ullevål Stadion, 0806 Oslo, Norway.

E-mail: anne.rustaden@inn.no (A.M. Rustaden).

Conclusions: Twelve weeks of BodyPump, heavy load resistance training with a personal trainer and non-supervised heavy load resistance training did not show any effect on self-reported musculoskeletal pain in overweight women.

Clinical Trial registration number: NCT01993953.

(<https://clinicaltrials.gov/ct2/show/NCT01993953>).

© 2019 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Being overweight (body mass index (BMI) ≥ 25.0 kg/m²) and obese (BMI ≥ 30.0 kg/m²) are common contributors to pain and disability in the musculoskeletal system,^{1–7} especially in the female population.³ With 40% of women worldwide classified as overweight⁸ and 15% as obese,⁹ the association between pain and BMI may evolve as an increasing challenge for the health care system.⁴

The pain-BMI association is primarily explained by increased mechanical load on joints and tissues, muscular inflammations and psychological issues.^{6,10,11} Consequently, individuals experience reduced physical ability and activity of daily living (ADL), as well as impaired quality of life.¹² In addition, overweight and obese women have previously reported fear of pain or injuries during exercise as a major barrier for a more active lifestyle, as well as an important deterrent for adherence to exercise.⁶ However, physical activity, and especially resistance training, may prevent or reverse pain symptoms by increasing muscle mass, muscle strength and physical function,^{12–14} help stabilizing the joints, improve mobility and improve proprioception.⁴

Resistance training can be practiced in many different ways, e.g. with a large range in repetitions and loads, and in different exercise settings (individually and in groups). In the current physical activity recommendations, overweight and obese individuals are recommended to perform resistance training 2–3 times/week, with an intensity between 60 and 80% of maximal muscle strength.¹⁵ However, high-repetition low-load resistance training sessions in groups, e.g. BodyPump, are popular exercise programs for women.¹⁶ Worldwide, BodyPump is distributed to over 14 000 fitness clubs, and weekly over 5 million individuals participate in a session.¹⁶ We have previously reported¹⁷ that 12 weeks of BodyPump was ineffective in improving muscle strength and body composition in overweight and obese women, compared to an inactive control group. In addition, we found that supervised heavy-load resistance training, in accordance with the American College of Sports Medicine (ACSM) recommendations, effectively improved muscle strength, and provided significantly higher exercise adherence, compared to non-supervised resistance training and BodyPump.¹⁷ However, we have not been able to find studies examining whether popular resistance training modalities, available in health- and fitness clubs, affect self-reported pain in overweight or obese women. Hence, the aim of this secondary analysis was to investigate the effect of three different resistance training modalities,

compared to controls, on musculoskeletal pain, in overweight and obese women. Furthermore, we aimed to study whether the results were influenced by adherence to exercise.

Material and methods

Study design

This is secondary analysis of a four armed assessor blinded randomized controlled trial (RCT).¹⁷ In the present study, primary outcome was to examine between group differences in musculoskeletal pain in previously inactive women with BMI ≥ 25.0 , allocated to twelve weeks of either BodyPump, heavy load resistance training with a personal trainer, non-supervised heavy load resistance training or no exercise. Secondarily, we examined whether there were differences in musculoskeletal pain between those reporting low versus high exercise adherence.

Study population

Participants were recruited via various social media channels and the homepage of the Norwegian School of Sport Sciences (NSSS). In total, 195 women contacted the principal investigator by phone or email. After aims and implications of the study were explained, eligibility criteria checked and a check-off health-profile scheme including health issues contraindicated for participation was fulfilled, a final sample of 143 participants were included. These participants were randomly allocated with concealed envelopes to either BodyPump (BP) ($n = 37$), heavy load resistance training with a personal trainer (PT) ($n = 35$), non-supervised heavy load resistance training (NS) ($n = 35$) or a non-exercising control group (C) ($n = 36$). An independent statistician performed block randomization, using a computer generated random numbers and an 8-persons block size, meaning that for every eight randomized person each block had two participants with the same intervention. The first 140 included participants were randomized with $n = 35$ in each group. Then, three more participants were included, randomized from a new 8-person block, giving different n in the four groups. Eligibility criteria's included BMI ≥ 25.0 , ages between 18 and 65 years and not regularly exercising defined as "not performing regular structured exercise $\geq$ twice a week the last six months".¹⁸ Using a inclusion screening scheme, participants were excluded if they had a history

Table 1 Exercise program BodyPump.

Music nr.	Exercise	Volume (Reps)
1 Warming-up	Straight leg deadlift, rowing, shoulder press, squat, lunges and bicepscurl	88
2 Leg	Squat	95
3 Chest	Bench press	80
4 Back	Rowing, stiff legged deadlift, clean & press and power press	75
5 Triceps	French press, tricepspress, pullover and overhead tricepspress	78
6 Biceps	Bicepscurl	68
7 Leg	Squat, lunges and squat jump	72 + 24 jumps
8 Shoulders	Push up, lateral raise, rowing and shoulderpress	76 + 36 push up
9 Stomach	Sit-ups, sit-ups to the side and side-plank	51 + 30 s

of diseases or injuries being contraindicated for the assessments or intervention (e.g. low back pain with radiation or osteoarthritis during the last six months, osteoporosis, secondary hypertension, history of coronary heart disease, stroke, arrhythmias, diabetes type 1 and neurological diseases). In addition, planned vacation or absence from exercise during the intervention period for >2 weeks, pregnancy, obesity surgery or psychiatric diseases were exclusion criteria's.

The study was prospectively registered in the Clinical Trial.gov Protocol Registration System (NCT01993953) and was approved by the Regional Committee for Medical Research Ethics Norway, Oslo (REK 2012/783). The procedures followed the World Medical Association Declaration of Helsinki, and all participants signed a written consent statement before entering the study.

Procedures and interventions

The participants were prescribed three exercise sessions weekly for twelve weeks, and the exercises were performed in a real life setting. Participants in the BP group had free access to several health- and fitness clubs offering BodyPump during the intervention period, while participants in the PT and NS group exercised at the health- and fitness club at the university.

BodyPump is a pre-choreographed, strengthening workout session, guided by a LesMills certified instructor. Every third month Les Mills releases a new program, but they are all based on the same principles and have the same structure.¹⁶ During the intervention period, BodyPump release no.83 was present at all fitness clubs (Table 1). Each BodyPump session consists of ten music tracks, 4–6 min each, including strengthening exercises targeting specific muscle groups. The participants exercise with a weight bar (1.25 kg), plates (1, 2.5 or 5 kg) and a step. In total, each 1-h session includes between 800 and 1000 repetitions, in the range of 50–100 repetitions in each muscle group. There are 1–2 min rest periods between each track, used to change weights and prepare for the next exercises. Training loads were self-selected, based on technique and intensity guidance from the instructor, as well as experiences from previous exercise sessions.¹⁶

The PT- and NS group followed a standardized nonlinear periodization program, including similar exercises as the BodyPump program. However, the number of repetitions

and sets continuously varied; 8–10 repetitions in session one, 13–15 repetitions in session two and 3–6 repetitions in session three. In week 1–4 the participants performed two sets in all the exercise, in week 5–8 they performed 2–3 sets, and in week 9–12 they increased to 3–4 sets. The rest periods varied between 60 and 120 s, with longest rest period in the heaviest sessions. At the beginning of each exercise session, the participants performed a 5–10 min low intensity warm-up on a treadmill or cycle ergometer. Details of the program have been previously reported.¹⁷ The participants were instructed to perform repetition maximum (RM) in each set, with proper lifting technique. Sixteen personal trainers took part in the study, all educated with a bachelor degree in physical activity and health, including a personal trainer certificate from our university. Participants in the PT group exercised together with their personal trainer in all sessions, and received continuously advice on appropriate training loads and lifting technique, as well as support and motivation during exercise. Participants in the NS group exercised on their own, except of one introduction session with a personal trainer who introduced them to the exercise program (proper lifting technique, training loads and progression), and a follow-up session after six weeks of exercise.

Participants in the C-group were asked to continue their usual lifestyle and ADL. If they performed any exercise or physical activity, they were asked to report this in a similar training diary as the intervention groups. After post-test, all controls were offered BodyPump classes for twelve weeks, and one resistance training session with a personal trainer.

Assessments

Musculoskeletal pain

Musculoskeletal pain was registered using the Standardized Nordic Pain Questionnaire (SNQ), developed to measure the prevalence of musculoskeletal pain and syndromes in epidemiological studies.¹⁹ The questionnaire registers whether the participants have experienced musculoskeletal pain in ten different anatomical body parts, during the last twelve months, and the last seven days. The participants self-reported the questionnaire at baseline (before randomization) with a blinded assessor present. At post-test, the same procedure was followed, but this time the participants were not blinded. For the purpose of the present study,




Figure 1 Flow chart of participants throughout the study.

responses to the last seven days were used as the primary outcome. In addition, responses the last twelve months were used in the descriptive analysis at baseline. The ten anatomical body parts included in the questionnaire were; head, neck, shoulder, elbow, wrist, low back, upper back, hip, knee and feet. All questions were formulated as e.g. "Have you ever during the last 12 months/7 days experienced pain in the...?". Possible responses were yes or no, and those who answered "yes", were categorized as having pain. The SNQ questionnaire have demonstrated adequate test-retest reliability (0–23% variation), and has previously been validated against clinical history with a variation between 0 and 20%.¹⁹

Adherence

The participants self-registered adherence to exercise in a training diary. For the purpose of this study, high adherence was defined as $\geq 75\%$ attendance to exercise (≥ 28 sessions of 36 possible), as this represented minimum two exercise sessions weekly. Low adherence was set at $\leq 75\%$ (≤ 27 sessions of 36 possible).

Statistical analysis

Analyzes were done with SPSS statistics program, version 21 (IBM Corporation, Route, Somers, NY, USA). Background data are presented as means with standard deviations (SD), and data on self-reported pain is presented as numbers (n) with percentages (%). One-way ANOVA was used to analyze possible differences between the groups in background variables and primary outcome. An attrition rate analysis of baseline characteristics between completers and non-completers was made with an independent t-test. Because of a high drop-out and low exercise adherence among our participants, results are presented for completers only (i.e. per-protocol analysis). McNemar's test was used to analyze if there was a difference in the proportion of the participants (the three intervention groups collapsed together) reporting muscle pain in any of the body parts prior to, versus

after the intervention. Chi-square test was used to analyze differences between the groups in self-reported pain (categorical data), as well as the differences in pain (yes/no) and high/low adherence. The percent difference between the groups was estimated with a 95% Wald confidence interval with Bonett–Price adjustment. Level of statistical significance was set at $p < 0.05$.

Results

Of 143 women randomized, 92 completed the study (mean age $39 \text{ years} \pm 10$, BMI $31 \pm 5 \text{ kg/m}^2$), with the following distribution: BP ($n = 24$, 65%), PT ($n = 28$, 80%), NS ($n = 19$, 54%) and C ($n = 21$, 58%) group (Fig. 1).

No statistically significant differences were found between the four groups in background data or musculoskeletal pain (Table 2), nor in the attrition rate analysis between completers and non-completers. Of totally 36 exercise sessions prescribed, adherence to exercise were 54% (± 20) in the BP group, 83% (± 15) in the PT group and 69% (± 20) in the NS group. The adherence was significantly higher in the PT group compared to the BP group ($29 \pm 5\%$, $p \leq 0.001$, 95% CI 15.5–41.5) and the NS group ($13.8 \pm 5\%$, $p = 0.031$, 95% CI 1.0–26.7), and in the NS group compared to the BP group ($14.7 \pm 6\%$, $p = 0.041$, 95% CI 0.5–28.9).

There were no between-group differences in musculoskeletal pain in any of the body parts at post-test (Table 3). Therefore, the prevalence of pain in the different body parts in the three intervention groups are collapsed and presented in Table 4. The analysis showed no statistical significant changes in reported pain from baseline to post-test. At baseline, the body parts with the highest reported pain was the neck, head, shoulder and lower back. At post-test, the highest prevalence was reported in the neck, shoulder, head and knees (Table 4).

Sub analyses of participants divided in high ($n = 38$) and low ($n = 20$) exercise adherence and report of musculoskeletal pain, is presented in Table 5. Irrespective of group allocation, there were no statistically significant difference in prevalence of bodily pain between those with high versus low adherence to exercise.

Table 2 Demographic data of the participants in the BodyPump group (BP), personal trainer group (PT), non-supervised group (NS) and control group (C).

Variable	BP (n = 24)	PT (n = 28)	NS (n = 19)	C (n = 21)
Age (year)	38 (11)	37 (9)	42 (12)	40 (12)
Weight (kg)	83 (11)	94 (21)	85 (14)	87 (15)
Height (cm)	168 (6)	169 (6)	168 (6)	167 (4)
BMI (kg/m ²)	30 (4)	33 (6)	30 (5)	31 (5)
Children (yes)	21	15	18	20
Daily smoker	9	2	4	4

Discussion

The present study investigated the effect of musculoskeletal pain in untrained women with BMI ≥ 25.0 , after twelve weeks of either BodyPump, heavy load resistance training with a personal trainer or non-supervised heavy load resistance training. None of the exercise modalities had effect on musculoskeletal pain, compared to inactive controls, and there were no difference between those with low and high adherence.

The present study included healthy women, and participants with a history of diseases or injuries contraindicated for resistance training were excluded. Hence, we assumed a low to moderate prevalence of musculoskeletal pain at baseline. However, 57 of the 143 participants reported pain in the SQN at baseline, in one or more body parts, with the head, neck, shoulder and lower back being the four most affected body parts. One reason for this, may be that the inclusion screening scheme contained specific diagnoses only (such as low back pain with radiation, osteoarthritis, osteoporosis, secondary hypertension, history of coronary heart disease, stroke, arrhythmias, diabetes type 1 and neurological diseases). Thus, musculoskeletal pain reported in the SQN at baseline, was not considered to be too severe by either the researcher or the participants in the study. Importantly, there were no group differences in self-reported pain in the SQN at baseline, in any of the body parts. In addition, the number of participants reporting pain in one or more body parts in the present study are comparable with prevalence of musculoskeletal pain in the general adult population.^{20,21} Cimmino et al.²⁰ report that 14–47% of the general population are affected of musculoskeletal pain at any given time, and Stone and Broderick²¹ report that obese individuals have 20% more daily pain than normal-weighted individuals. In addition, the most affected body parts in our study group have all previously been reported with high risk of pain in overweight and obese individuals.^{22–25} However, comparison of pain between studies is difficult, due to differences in the definition used, study population and measurement methods.²⁶

The overweight-pain association is related to factors as a sedentary lifestyle, reduced physical function and low muscle strength.^{2,4,27} Interestingly, several studies have found that resistance training effectively reverse or improve pain symptoms in overweight individuals,^{1,2,5,6} particularly low-back pain.^{2,13,14} In the present study, low back pain was non-significantly reduced with 11% from baseline to

post-test. Lack of power, low back pain with radiation as one of the exclusion criteria's, low responsiveness of the questionnaire or the content of the program, which had no special focus on treating low back pain, might be explanations why our reduction was somewhat smaller than in other studies.

Unfortunately, the present study had a high drop-out rate. As Herbert et al.²⁸ state that more than 85% of the participants should be assessed and obtained in the main outcome when implementing intention-to treat analysis (ITT),²⁸ we used completers only in the present analysis. In total, seven participants from the three intervention groups dropped out because of illness/injury (Fig. 1). We do not know whether these were exercise-induced injuries, as most of the participants did not give any reason for drop-out. As much as 20 participants were lost to follow-up without reporting reason, the majority of these from the NS group. In addition, the exercise adherence was low in the BP and NS group, which may have affected the outcomes. However, the results did not change when we compared participants with high and low adherence. High drop-out from exercise and low exercise adherence are well-known challenges in overweight and obese individuals.^{2,6} Interestingly, Zdziarski et al.² emphasized that exercise modifications, as low load as an alternative of high load resistance training, could reduce acute exercise induced pain, and possibly increase exercise adherence among obese individuals.² This was not observed in the present study, as the BP group, representing low load resistance training, had significantly lower exercise adherence, compared to the two heavy load groups (PT and NS). However, the fact that our participants exercising with a personal trainer had higher adherence, compared to non-supervised exercise, corresponds with previous findings.⁶ Arikawa et al.⁶ compared supervised (month 1–4) and non-supervised (month 5–24) resistance training in untrained overweight and obese women, and found significantly higher adherence during the supervised period.⁶ Thus, support and motivation from certified personal trainers might be a key-factor to avoid drop-out and increase exercise adherence in this study group.⁶

Stidsen et al.²⁹ investigated musculoskeletal issues among fitness club members, using the same questionnaire as the present study (SQN). They found that 56% of totally 500 new fitness club members reported musculoskeletal pain in one or more body parts when joining the club, and 77% of these stated that pain was one of the main reasons why they joined the club.²⁹ Thus, it is important

Table 3 Self-reported musculoskeletal pain at baseline and post-test in the BodyPump group (BP), personal trainer group (PT), non-supervised group (NS) and control group (C). Differences between the groups analyzed with chi-square test, presented with *p*-value. Presented as numbers reporting pain (*n*)/of the total number of participants (*n*) in each group, and as percent (%).

Body part	Baseline BP <i>n</i> (%)	PT <i>n</i> (%)	NS <i>n</i> (%)	C <i>n</i> (%)	<i>p</i> -Value	Post-test BP <i>n</i> (%)	PT <i>n</i> (%)	NS <i>n</i> (%)	C <i>n</i> (%)	<i>p</i> -Value
Head	8/31 (26%)	10/32 (31%)	8/25 (32%)	8/31 (26%)	0.92	8/24 (33%)	7/28 (25%)	1/19 (5%)	8/21 (38%)	0.09
Neck	10/31 (32%)	8/32 (25%)	13/27 (48%)	12/31 (39%)	0.30	11/24 (46%)	6/28 (21%)	5/19 (26%)	4/21 (19%)	0.16
Shoulder	8/31 (26%)	9/32 (28%)	7/28 (27%)	11/31 (35%)	0.84	6/24 (25%)	9/28 (32%)	6/19 (32%)	4/21 (19%)	0.73
Elbow	1/31 (3%)	1/32 (3%)	4/25 (16%)	2/30 (7%)	0.20	0/24 (0%)	4/28 (14%)	5/19 (26%)	2/21 (10%)	0.07
Wrist	2/30 (7%)	3/32 (9%)	2/25 (8%)	3/30 (10%)	0.97	4/24 (17%)	4/28 (14%)	0/19 (0%)	2/21 (10%)	0.32
Upper back	4/30 (13%)	6/32 (19%)	4/25 (16%)	4/30 (13%)	0.92	0/24 (0%)	5/28 (18%)	2/19 (11%)	2/21 (10%)	0.20
Lower back	8/31 (26%)	7/32 (22%)	9/27 (33%)	10/30 (33%)	0.96	2/24 (8%)	5/28 (18%)	4/19 (21%)	3/21 (14%)	0.67
Hip	5/31 (16%)	2/32 (6%)	2/25 (8%)	3/30 (10%)	0.60	0/24 (0%)	3/28 (11%)	0/19 (0%)	1/21 (5%)	0.20
Knee	2/34 (6%)	5/32 (16%)	6/25 (24%)	2/30 (7%)	0.13	3/24 (13%)	8/28 (29%)	5/19 (26%)	2/21 (10%)	0.25
Feet	2/32 (6%)	5/32 (16%)	3/25 (12%)	5/30 (17%)	0.59	2/24 (8%)	4/27 (15%)	2/19 (11%)	4/21 (19%)	0.73

Table 4 The number of participants that reported musculoskeletal pain in the different body parts at baseline and post-test, with the three intervention groups collapsed. Reported as number participants reporting pain (*n*)/total number of participants (*n*), and percent (%). Differences from baseline to post-test, analyzed with McNemar's test are presented with *p*-value, and percent group differences (%) estimated with a 95% Wald confidence interval with Bonett-Price adjustment.

Body part	Baseline n/n total (%)	Post-test n/n total (%)	<i>p</i> -Value	Percent diff. (95% CI)
Head	26/88 (29%)	16/71 (23%)	1.00	1.5% (−9.55 to 12.45)
Neck	31/90 (34%)	22/71 (31%)	1.00	0% (−11.88 to 11.88)
Shoulder	24/89 (27%)	21/71 (30%)	0.66	−4.5% (−17.93 to 9.24)
Elbow	6/88 (7%)	9/71 (13%)	0.06	−7.6% (−14.99 to 0.07)
Wrist	7/87 (8%)	8/71 (11%)	1.00	0% (−10.3 to 10.3)
Upper back	14/87 (16%)	7/71 (10%)	0.23	7.7% (−2.93 to 17.86)
Lower back	24/90 (27%)	11/71 (16%)	0.18	−1.2% (−12.23 to 9.88)
Hip	9/88 (10%)	3/71 (4%)	1.00	0% (−7.17 to 7.17)
Knee	13/91 (14%)	16/71 (23%)	0.09	−10.4% (−20.88 to 0.59)
Feet	10/89 (11%)	8/70 (11%)	1.00	−1.5% (−12.03 to 9.05)

Table 5 Differences in self-reported musculoskeletal pain (yes/no) and high ($\geq 75\%$) versus low ($\leq 75\%$) exercise adherence. Analyzed with chi-square test, and presented with *p*-value. Differences between the groups estimated with a 95% Wald confidence interval with Bonett-Price adjustment, presented with percent % and 95% CI.

Body part	Yes/no	Low adherence <i>n</i> (%)	High adherence <i>n</i> (%)	<i>p</i> -Value	Percent diff. (95% CI)
Head	Yes No	6 (23%) 20 (77%)	6 (19%) 26 (81%)	0.69	24.2% (−7.09 to 39.58)
Neck	Yes No	8 (31%) 18 (69%)	9 (28%) 23 (72%)	0.83	15.5% (−2.18 to 32.18)
Shoulder	Yes No	9 (35%) 17 (65%)	10 (31%) 22 (69%)	0.79	12.1% (−5.68 to 29.01)
Elbow	Yes No	3 (12%) 23 (88%)	5 (16%) 27 (85%)	0.65	31.0% (−13.80 to 46.20)
Wrist	Yes No	2 (8%) 24 (92%)	5 (16%) 27 (84%)	0.36	32.7% (−15.34 to 47.99)
Upper back	Yes No	1 (4%) 25 (96%)	6 (19%) 26 (81%)	0.08	32.7% (−14.70 to 48.64)
Lower back	Yes No	4 (15%) 22 (45%)	5 (16%) 27 (84%)	0.98	29.3% (−12.27 to 44.40)
Hip	Yes No	0 (0%) 26 (100%)	2 (6%) 30 (94%)	0.20	41.4% (−25.25 to 54.75)
Knee	Yes No	3 (12%) 23 (88%)	9 (28%) 23 (72%)	0.12	24.1% (−5.22 to 41.44)
Feet	Yes No	2 (8%) 24 (92%)	3 (10%) 28 (90%)	0.79	36.8% (−20.18 to 51.01)

Assessed for eligibility *n* = 195.

to investigate the effect of pain during and after popular exercise modalities available in health- and fitness clubs, especially in risk groups like overweight and obese. Therefore, even though none of the three exercise modalities in the present study significantly reduced self-reported pain after 12 weeks of exercise, we find it important to highlight that no adverse effect was seen. Hence, the resistance training modalities in the present study may all be appropriate for overweight women. This knowledge is important to e.g. physical therapists aiming to include or recommend resistance training among overweight or obese patients. However, it is important to emphasize that resistance training may induce immediate exercise induced pain and give temporary delayed onset muscle soreness (DOMS).³⁰

Strengths of the present study were use of a randomized controlled design, blinded investigator and use of a validated questionnaire. The study had a high external and ecological validity, as the exercise training was performed in a real-life setting, under pragmatic conditions. In addition,

all participants followed the same standardized exercise programs, and the personal trainers followed the same standardized instructions. Limitations in the study are the high drop-out and the low exercise adherence in the BP and NS group. In addition, the SNQ do not distinguish between intensity and type of pain. Therefore, the participants may have interpreted the definition of musculoskeletal pain differently, and e.g. confused stiffness, DOMS, fatigue and functional limitations. Questionnaires including pain intensity e.g. the visual analog scale for pain, numeric rating scale for pain, McGill pain questionnaire or the short form 36 bodily pain scale,³¹ could have given more detailed information.

To conclude, our study showed no between- or within group changes in self-reported musculoskeletal pain after twelve weeks of either BodyPump (high-repetition low load resistance training), heavy load resistance training with a personal trainer or non-supervised heavy load resistance training.

Funding/support statement

The present study did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

Acknowledgements

We thank all the participating women, and Ingar Holme and Morten Fagerland for statistical advice.

References

- Kulkarni K, Karssiens T, Kumar V, Pandit H. Obesity and osteoarthritis. *Maturitas*. 2016;89:22–28.
- Zdziarski LA, Wasser JG, Vincent HK. Chronic pain management in the obese patient: a focused review of key challenges and potential exercise solutions. *J Pain Res*. 2015;8:63–77.
- Yoo JJ, Cho NH, Lim SH, Kim HA. Relationships between body mass index, fat mass, muscle mass, and musculoskeletal pain in community residents. *Arthritis Rheumatol*. 2014;66:3511–3520.
- Vincent HK, Adams MCB, Vincent KR, Hurley RW. Musculoskeletal pain, fear avoidance behaviors, and functional decline in obesity. *Reg Anesth Pain Med*. 2013;38:481–491.
- Kulie T, Slattengren A, Redmer J, Counts H, Eglash A, Schrager S. Obesity and women's health: an evidence-based review. *J Am Board Fam Med*. 2011;24:75–85.
- Anandacoomarasamy A, Caterson I, Sambrook P, Fransen M, March L. The impact of obesity on the musculoskeletal system. *Int J Obes*. 2008;32:211–222.
- Urek R, Crncević-Urek M, Cubrilo-Turek M. Obesity – a global public health problem. *Acta Med Croat*. 2007;61: 161–164.
- Ng M, Fleming T, Robinson M, et al. Global, regional and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease study 2013. *Lancet*. 2014;384: 766–781.
- Di Cesare M, Bentham J, Stevens GA, et al. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. 2016;387:1377–1396.
- Clark JE, Goon DT. The role of resistance training for treatment of obesity related health issues and for changing health status of the individual who is overfat or obese: a review. *J Sports Med Phys Fitness*. 2015;55:205–222.
- Ray L, Lipton RB, Zimmerman ME, Katz MJ, Derby CA. Mechanisms of association between obesity and chronic pain in the elderly. *Pain*. 2011;52:53–59.
- Barry VW, Baruth M, Beets MW, Durstine JL, Liu J, Blair SN. Fitness vs. fatness on all-cause mortality: a meta-analysis. *Prog Cardiovasc Dis*. 2014;56:382–390.
- Wasser JG, Vasilopoulos T, Zdziarski LA, Vincent HK. Exercise benefits for chronic low back pain in overweight and obese individuals. *Res Sports Med*. 2017;9:181–192.
- Vincent HK, Vincent KR, Seay AN, Conrad BP, Hurley RW, George SZ. Back strength predicts walking improvement in obese, older adults with chronic low back pain. *PM R*. 2014;6:418–426.
- Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. American College of Sports Medicine Position Stand: appropriate physical activity intervention on strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc*. 2009;41:459–471.
- Les Mills International Web site. Available at: <http://www.lesmills.com/> Accessed 15.08.18.
- Rustaden AM, Haakstad LAH, Paulsen G, Bø K. Effects of Body-Pump and resistance training with and without a personal trainer on muscle strength and body composition in overweight and obese women – a randomised controlled trial. *Obes Res Clin Pract*. 2017;11:728–739.
- Haakstad LA, Bø K. Effect of a regular exercise programme on pelvis girdle and low back pain in previously inactive pregnant women: a randomized controlled trial. *J Rehabil Med*. 2015;47:229–234.
- Kuorinka I, Jonsson B, Kilbom A, et al. Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Appl Ergon*. 1987;18:233–237.
- Cimmino MA, Ferrone C, Cutolo M. Epidemiology of chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol*. 2011;25:173–183.
- Stone AA, Broderick JE. Obesity and pain are associated in the United States. *Obesity*. 2012;20:1491–1495.
- Pavlovic JM, Vieira JR, Lipton RB, Bond DS. Association between obesity and migraine in women. *Curr Pain Headache Rep*. 2017;21:41.
- Zhang TT, Liu Z, Liu YL, Zhao JJ, Liu DW, Tian QB. Obesity as a risk factor for low back pain: a meta-analysis. *Clin Spine Surg*. 2018;31:22–27.
- Chai NC, Scher AI, Moghekar A, Bond DS, Peterlin BL. Obesity and headache: Part 1 – Systematic review of the epidemiology of obesity and headache. *Headache*. 2014;54:219–234.
- Lund Nilsen TI, Holtermann A, Mork PJ. Physical exercise, body mass index, and risk of chronic pain in the low back and neck/shoulders: longitudinal data from the Nord-Trøndelag health study. *Am J Epidemiol*. 2011;174:267–273.
- Janke EA, Collins A, Kozak AT. Overview of the relationship between pain and obesity: what do we know? Where do we go next? *J Rehabil Res Dev*. 2007;44:245–326.
- Blümel JE, Arteaga E, Mezones-Holguín E, Zúñiga MC, Witis S, Vallejo MS. Obesity is associated with a higher prevalence of musculoskeletal pain in middle-aged women. *Gynecol Endocrinol*. 2017;33:378–382.
- Herbert R, Jamtvedt G, Mead J, Hagen KB. *Physical Evidence-Based Physiotherapy*. Butterworth Heinemann: Elsevier; 2005.
- Stidsen L, Jensen J, Kvorning T, Hartvigsen J. Are fitness centres part of the health care sector? *Ugeskr Laeger*. 2008;170:2790–2794.
- Dannecker EA, Koltyn KF. Pain during and within hours after exercise in healthy adults. *Sports Med*. 2014;44:921–942.
- Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain. *Arthritis Care Res*. 2011;63:240–252.