

## MASTERCLASS

# Culture and musculoskeletal pain: strategies, challenges, and future directions to develop culturally sensitive physical therapy care



Felipe J.J. Reis<sup>a,b,c,\*</sup>, Jo Nijs<sup>c,d,e</sup>, Romy Parker<sup>f</sup>, Saurab Sharma, PhD<sup>g,h</sup>, Timothy H. Wideman<sup>i</sup>

<sup>a</sup> Physical Therapy Department, Instituto Federal do Rio de Janeiro (IFRJ), Rio de Janeiro, Brazil

<sup>b</sup> Postgraduation Program, Clinical Medicine Department, Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>c</sup> Pain in Motion Research Group, Department of Physical Therapy, Human Physiology and Anatomy, Faculty of Physical Education & Physical Therapy, Vrije Universiteit Brussel, Brussels, Belgium

<sup>d</sup> Department of Physical Medicine and Physical Therapy, University Hospital Brussels, Brussels, Belgium

<sup>e</sup> Unit of Physical Therapy, Department of Health and Rehabilitation, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

<sup>f</sup> Department of Anaesthesia and Perioperative Medicine, Groote Schuur Hospital, University of Cape Town, Anzio Rd, Observatory, Cape Town, Western Cape, South Africa

<sup>g</sup> Department of Exercise Physiology, School of Health Sciences, Faculty of Medicine and Health, University of New South Wales, Sydney, Australia

<sup>h</sup> Centre for Pain IMPACT, Neuroscience Research Australia, Sydney, Australia

<sup>i</sup> School of Physical and Occupational Therapy, McGill University, Montreal, Canada

Received 6 October 2021; received in revised form 13 July 2022; accepted 25 August 2022

Available online 15 September 2022

### KEYWORDS

Chronic pain;  
Culture;  
Exercise;  
Pain;  
Patient education as topic

### Abstract

**Background:** Pain experience has a multidimensional nature. Assessment and treatment recommendations for pain conditions suggest clinicians use biopsychosocial approaches to treat pain and disability. The current pain research is overwhelmingly skewed towards the study of biological and psychological factors including interventions, whereas, cultural factors are often ignored.

**Objective:** The aims of this Masterclass is threefold: (1) to discuss cultural influences on pain, (2) to provide strategies for delivering appropriate pain education and exercises in culturally diverse people with chronic pain, and (3) to present challenges and future directions to clinicians and researchers.

**Discussion:** Cultural factors have a relevant influence on the way individuals experience and manage health and illness. Thus, people with different cultural experience perceive, respond, communicate and manage their pain in different ways. In this aspect, the contents of pain

\* Corresponding author at: Instituto Federal do Rio de Janeiro, Campus Realengo, Rua Carlos Wenceslau, 343, Realengo. CEP 21715-000, Rio de Janeiro, RJ, Brazil.

E-mail: [felipe.reis@ifrj.edu.br](mailto:felipe.reis@ifrj.edu.br) (F.J. Reis).

education should be presented using different culturally appropriate examples, metaphors, images, and delivery methods that may enhance the impact of the message. Efforts should be made to produce and spread culturally adapted evidence-based materials and resources. In addition, a culturally sensitive approach may help to introduce patients to graded activities, so that they can apply these strategies in culturally acceptable and meaningful ways. Future studies should investigate the effectiveness of culturally-adapted interventions in pain-related outcomes in different pain conditions in patients with different cultural backgrounds.

© 2022 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier España, S.L.U. All rights reserved.

## Introduction

Chronic musculoskeletal pain is a global health issue.<sup>1</sup> Current evidence indicates that pain is modulated by biological, emotional, cognitive, behavioral, environmental, and socio-cultural factors.<sup>2–4</sup> However, the majority of literature emphasizes the influence of biological and psychological aspects on pain, with the influence of the socio and cultural aspects of pain largely overlooked.<sup>5–8</sup>

The literature focusing on the influential role of culture on pain has considered variables such as ethnicity, gender, nationality, and language as proxy measures of culture as these variables may reflect cultural differences.<sup>9,10</sup> However, according to The United Nations Educational, Scientific and Cultural Organization (UNESCO), culture is defined as “the set of distinctive spiritual, material, intellectual and emotional features of society or a social group, and that it encompasses, in addition to art and literature, lifestyles, ways of living together, value systems, traditions and beliefs”.<sup>11</sup> Therefore, an individual can identify themselves as belonging to more than one social group, based on ethnicity, gender, nationality, common interests, or sexual orientation.<sup>12</sup> This intersectionality of belonging makes the culture of an individual dynamic and constantly evolving.

Cultural factors have a relevant influence on the way individuals experience and manage health and illness, as well as receptivity to health interventions, their pain behaviors (e.g., coping responses, help seeking behavior), and beliefs (e.g., causes of pain, consequences, how controllable pain is).<sup>13</sup> Beliefs, pain-related fear, catastrophizing, and avoidance are key factors in the development of disability in chronic musculoskeletal pain.<sup>13,14</sup> Pain catastrophizing, an important component of the fear-avoidance model, has been consistently regarded as one of the most influential determinants of pain-related outcomes.<sup>15</sup> The fear-avoidance model suggests that pain may be interpreted as threatening bodily integrity, which may lead to people prioritizing pain control over achieving valued life goals,<sup>16</sup> leading to a vicious cycle involving catastrophizing, hypervigilance, and avoidance or escape behavior and, in turn, leading to disuse, disability, depression and increased pain.<sup>16</sup> Although pain catastrophizing can be considered as a maladaptive belief (i.e., negative interpretation about the severity and the concept that unexplained symptoms are usually serious),<sup>17</sup> this negative cognitive appraisal can be related to other reasons in different cultures (e.g., the feeling of abandonment by God during their suffering).<sup>18</sup>

Although some attempts have been made to explore the influence of cultural factors on pain, pain catastrophizing, beliefs, and coping strategies, it remains challenging for

clinicians to use this information in their practice.<sup>19</sup> For instance, while clinical practice guidelines recommend addressing misconceptions about pain as one of the first line of interventions in patients with musculoskeletal pain,<sup>20</sup> the implementation of such interventions in culturally and linguistically diverse communities remain challenging in clinical practice.<sup>21</sup> Given these considerations, the aims of this Masterclass are: (i) to discuss cultural influences on pain, pain catastrophizing, pain beliefs and behavior, (ii) to provide strategies to deliver pain education and exercises in culturally diverse people with chronic musculoskeletal pain, and (iii) to present challenges and future directions to clinicians and researchers about culture and pain.

## How can culture influence pain, beliefs, and pain catastrophizing?

The literature highlights the role of race/ethnicity<sup>22,23</sup> and sex/gender<sup>24</sup> differences in pain prevalence, coping, sensitivity, and tolerance. For example, African Americans, Asians, and Hispanics experience and report more intense pain compared to their non-Hispanic White counterparts.<sup>23</sup> Despite this, African Americans and other ethnic minorities receive lesser and poorer care for their pain.<sup>25</sup> Sex/gender differences can also be related to cultural factors. In some cultures, boys and men are taught to tolerate pain, and endure painful experiences, while girls and women are taught to be sensitive, careful, and to verbalize discomfort.<sup>26</sup>

Cultural variability in beliefs regarding the causes and consequences of pain play a significant role in the patient's pain experience.<sup>13,27</sup> For example, in Mexican-American culture, pain can be viewed as being due to God's will, punishment or penance.<sup>27</sup> The Aboriginal Australians with chronic low back pain did not express their pain and demonstrated few pain behaviors because of cultural beliefs that pain is a social and spiritual dysfunction.<sup>28</sup> In Zulu culture, pain may be considered a consequence of bewitching.<sup>29</sup> Family beliefs (e.g., enduring pain, the role of folk healers), spirituality, and religiosity<sup>30</sup> are suggested to play a role in how pain is expressed and managed in Hispanic cultures.<sup>31</sup> In case of beliefs emphasizing supernatural or divine causes of pain, the external locus of control can be associated with increased pain intensity.<sup>32</sup>

Evidence suggests that cultural beliefs can be changed following healthcare interactions.<sup>33</sup> For example, people from rural Nepal tend to perceive back pain as a “normal” part of aging and they do not seek care;<sup>34,35</sup> whereas, back

pain is considered a biological phenomenon in urban Nepal,<sup>36</sup> resulting in greater healthcare utilization. Aboriginal Australians with chronic low back pain presented negative beliefs and misconceptions about their condition among those who were exposed to healthcare interactions.<sup>33</sup> The influence of the dominant health approach in Western societies appeared to contribute to unhelpful pain beliefs as an example of acculturation (i.e., a process by which individuals adopt beliefs and behaviours of another dominant culture).<sup>33</sup>

People with different cultural experience perceive, respond, communicate and manage their pain in different ways.<sup>27</sup> In Western society, people with chronic pain may experience shame, embarrassment, and humiliation as a consequence of pain-related disability.<sup>37</sup> Sharma et al.<sup>19</sup> compared pain-related beliefs and catastrophizing across countries, language groups, and country economy. The authors found that pain-related beliefs were different in between-country and between-economic region comparison. However, there were no difference in pain beliefs in people living in the same country but speaking different languages. Pain catastrophizing was greater in people with chronic pain from the United States than Portugal, and people from Australia presented more pain catastrophizing than samples from Denmark and Brazil. The interactions between pain catastrophizing, pain sensitivity, and culture/ethnicity should also be considered. Fabian et al.<sup>38</sup> found that only situational catastrophizing varied by ethnicity, with African Americans reporting greater catastrophizing than Asian/Pacific Islanders and Caucasians. The authors found that situational catastrophizing significantly mediated pain intensity. Another study found that pain catastrophizing partially mediated race differences in pain tolerance and mediated sex differences in pain intensity.<sup>39</sup>

## Culture and the dynamic biopsychosocial model

The biopsychosocial model, that is well endorsed in the pain field, supports the notion that biological, psychological, societal, and lifestyle factors are interconnected and mutually reinforcing.<sup>40</sup> This model has been criticized because it has been often applied in a fragmented manner in clinical practice and research.<sup>6–8</sup> The narrow focus on some psychological dimensions (e.g., behavior and cognition) and minimal consideration of the social dimensions (including culture) perpetuate a reductionist view of the pain experience.<sup>6,7</sup> It is important to highlight that the influence of bio, psycho and social dimensions on health are not fixed, but rather interact with each other over time.<sup>41</sup> Lehman et al.<sup>41</sup> proposed that the biopsychosocial model should be viewed as a dynamic model of health constructed as a product of the reciprocal influences of biological, psychological, interpersonal, and macrosystem contextual dynamics that unfold over personal and historical time. This model considers interpersonal dynamics over the life span including direct contact with others, as well as the reverberating consequences of others' actions. The interpersonal dynamics of health can be explained using Bronfenbrenner's ecological framework and concepts of microsystem, mesosystem, exosystem, and macrosystem factors (Fig. 1). The *microsystem* factors include family members, work environments, peers,




Fig. 1 Bronfenbrenner's ecological framework.

and community health resources. The *mesosystem* relies on interactions among various microsystem factors (e.g. multiple family members and social groups). The *exosystem* refers to an indirect influence when the microsystem is affected by dynamics within their own microsystems (e.g. influence from spouses' employment setting). The *macrosystem* (referenced as contextual dynamics) includes shared culture, norms, policies, and values.

Clinicians should be mindful that all biological, psychological, and social factors may occur within contextual dynamics and are influenced by culture. Thus, we suggest that during the biopsychosocial assessment of patients with chronic musculoskeletal pain, clinicians should use open and reflective questions to capture patients' pain experience, pain beliefs, and coping strategies.<sup>4</sup> Further, clinicians should avoid making stereotypical cultural judgments based on cultural affiliations. Clinicians should be respectful, open, empathetic, authentic, sensitive, and caring throughout their clinical encounter. Therefore, the concept of a non-judgmental patient-centered view should be cultivated and derogatory labels (both with colleagues and patients), such as "catastrophizer", "avoider", or "malingerer" should be avoided.<sup>42</sup> Box 1 provides example questions that clinicians can use to obtain a culturally sensitive pain assessment.

## Culture-sensitive pain neuroscience education and exercise therapy

Although several interventions [e.g. cognitive functional therapy,<sup>43,44</sup> group cognitive behavioral treatment,<sup>45</sup> acceptance and commitment therapy, cognitive behavioral therapy,<sup>46</sup> mindfulness,<sup>47</sup> multimodal treatment<sup>48</sup>], have been described to address maladaptive pain beliefs, pain catastrophizing, fear

Box 1 Examples of questions to address pain beliefs.	
Meaning of health and disease	What does good health mean to you? What does disease mean to you?
Meaning of pain	How do you describe your pain? What does your pain mean to you? What words can you use to describe your pain? Why do you think it hurts? What do you think is the cause of your pain?
Explanations	Do you have any explanation for your pain? Why do you think that your pain is lasting for a long time?
Cognition and Emotions	Given that you have pain, how do you feel about it? Do you feel unfair, angry, frustrated, or anxious because of your pain? Do you have any fear related to your pain? If so, what is your fear? Do you have any worries about your pain? If so, what are your worries? Do you think it is unfair that you suffer from (chronic or persistent) pain?
Behavior and coping	What do you do when you are in pain? Which activities do you avoid due to pain? How do you cope with your pain?
Openness to learn or change	Do you want to learn more about pain?
Treatments	What kind of treatment do you think you need? Do you usually try treatments such as healers, prayers, or plant preparations to help you with your pain? Does it help? What treatments have you tried in the past? Did they work? Why?
Expectancies	What are the most important results you hope to receive from the treatment? Do you think your pain will get better?
Social support	Do the people around you notice when you are in pain? Who knows about your persistent pain problem, and who does not? Do you have family and friends who help you with your pain? Who helps you? Who, if anyone, in your family do you talk to about your pain? What do they know? What do you want them to know?

### Pain neuroscience education (PNE)

PNE aims to decrease the perceived threatening nature of the patients’ pain to facilitate engagement with physical activities which specifically target beliefs, pain-related fear, pain catastrophizing, and avoidance. The current evidence suggests that PNE can be effective for several pain-related outcomes including pain intensity, disability, pain catastrophizing, and kinesiophobia in the short- to medium-term in patients with chronic musculoskeletal pain.<sup>53</sup> The impact of PNE on patients’ cognition and fear-avoidance beliefs is limited to small to medium effect sizes,<sup>53</sup> illustrating that more is needed than education alone. A recent review showed positive results in favor of the addition of PNE to physical therapy treatment on disability and pain intensity in patients with chronic low back pain.<sup>54</sup>

Patients want to receive a clear and consistent message about their pain, treatment options, and self-management strategies.<sup>55</sup> A culturally sensitive PNE session should take into account that patients have an expectation that the provider will be a good communicator and will listen to concerns, recognizing the legitimacy of their pain, and respect their culture, values, and preference. The focus of a PNE session should switch from “clinician-centered teaching” to “patient-centered learning.” Hence, clinicians should avoid imposing their knowledge about pain biology over patients’ concepts, values, and beliefs. During a clinical encounter, clinicians should use open-ended questions to capture the patients’ beliefs about their pain. This strategy provides insight into the patient’s cultural meaning of pain, beliefs, and coping strategies. Within the context of “patient-centered learning,” it is expected that clinician and patient build a two-way street for interaction (Fig. 2).

A culturally sensitive PNE session is important because patients present different explanations for their pain. For example, people from rural communities in Nepal have contrasting beliefs about pain ranging from “pain as a normal part of life or aging and that they should live with it” to “pain is a result of bad deeds from the past lives or a result of black magic”.<sup>34</sup> Some Brazilians can consider their pain as a loss of faith or even a punishment from God leading them to seek prayers and rituals to receive forgiveness. In some South African cultures, pain may be interpreted as a punishment from ancestors. Thus, it is important to consider that




Fig. 2 Differences between “clinician-centered teaching” and “patient-centered learning”.

of movement, pain intensity, and disability, this paper focuses on treatment approaches such as pain neuroscience education (PNE),<sup>49</sup> cognition-targeted exercise therapy,<sup>50</sup> behavioral graded activity,<sup>51</sup> and graded exposure.<sup>52</sup>



**Box 2** Pain education messages adapted for South Africa as an example that pain education concepts should be adapted considering the patients' context and cultural background.

Pain and the extra-sensitive alarm system

**Pain and the Anthills**

In some parts of the county, there are anthills (a nest above the ground of clay). If you go to an anthill and you make a hole in it using a rock, then all the ants come out to fix the hole and they attack you. But, if you do that every day, after a while, if you just walk close to the anthill, and you stomp your foot, all the ants come out to attack you before you do anything to their nest.

**Pain and the Fogbanks**

In some parts of the country there are fogbanks coming from the sea. It looks like smoke. In these areas bushfires are common. Thus, if you see the fogbank coming in, people may respond as if it is smoke and fire - a disaster, when it is just a fogbank.

the same set of target key PNE concepts may not necessarily work in all cultural groups. Clinicians need to identify key educational concepts in a pain education program that may help to produce a more focused, culturally appropriate, and succinct program.

Clinicians should consider several audiovisual aids to deliver PNE. While slide presentations work for western cultures, they did not seem to work in the context of Nepal as patients had no previous experience with this approach. Printed postcards appeared to be a good alternative in the Nepali context.<sup>36</sup> In a South African context, printed materials which patients can take home and share with friends and family are regarded as valuable and may enhance the impact of the message.<sup>56</sup>

Clinicians should acknowledge, respect, and validate rather than confront patients' beliefs and cultural values. It is helpful for patients if clinicians respect their beliefs but offer alternative explanations and draw patients to reflect on their pain problems and explain contributors to pain such as sleep, activity levels, stress, and mood. These are often effective if patients or their family members can draw examples from their own experiences. If not, other examples from local context may be used here to demonstrate these relationships. In **Box 2** we present some examples of metaphors adapted for South African to explain pain concepts.

### **Culturally relevant behavioral graded activity, graded exercise therapy, graded exposure in vivo, or cognition-targeted exercise therapy**

A culturally sensitive PNE approach may help introduce patients to graded activities (i.e., behavioral graded activity, graded exercise therapy, graded exposure in vivo, or cognition-targeted exercise therapy), so that patients can apply these strategies in culturally acceptable and meaningful ways. Moreover, without applying the knowledge regarding pain into daily life, deep learning and long-term effects cannot be expected.

The available evidence suggests that patients with chronic spinal pain who have received PNE (implying that patients have increased their understanding of their pain problems, and the threatening nature of pain [flares] is reduced), respond better to exercise therapy.<sup>57</sup> In line with this, it seems rational that this also accounts for adopting a more active lifestyle,<sup>58</sup> but direct evidence supporting this notion

is currently lacking. In addition, this is not a linear process, and responses can vary substantially from patient to patient. The synergistic effect of PNE combined with such active approaches was illustrated in a trial with 120 patients with chronic spinal pain randomized to PNE plus cognition-targeted exercise therapy or back/neck school education plus general exercise therapy. The trial found hardly any clinically important changes after the education phase, but medium to large improvements were seen after the cognition-targeted exercise therapy on kinesiophobia and hypervigilance even at 1-year follow-up.<sup>57</sup> Both active approaches were associated with marked reductions in pain catastrophizing.<sup>57</sup>

One key aspect includes changing from the classical symptom-contingent approach ("Stop the activity or exercise once it hurts") to a time-contingent approach to exercises and daily activities ("Perform the activity or exercise for five minutes, regardless of the pain"). Behavioral graded activity can be used to implement such a time-contingent approach into the patients' daily life. Behavioral graded activity can be regarded as a culturally sensitive approach as it is a behavioral treatment integrating the concept of operant conditioning to increase the level of physical activity in the patient's daily life.<sup>59</sup> Behavioral graded activity is an effective approach for patients with chronic low back pain<sup>60</sup> and osteoarthritis.<sup>59</sup>

For addressing maladaptive beliefs and pain catastrophizing, especially for highly feared activities and exercises, other approaches are needed. Approaches that are useful for tackling such highly feared activities include cognition-targeted exercise therapy and graded exposure. The current evidence suggests that cognition-targeted exercise therapy presented superior results over pain-contingent exercise therapy in patients with chronic low back pain.<sup>57</sup> Limited evidence suggests that graded exposure is more effective than behavioral graded activity for improving disability and catastrophizing at short term in patients with chronic low back pain.<sup>60</sup> Graded exposure in vivo is a cognitive behavioral treatment approach that has resulted in positive outcomes in patients with chronic low back pain,<sup>61</sup> complex regional pain syndrome,<sup>62</sup> whiplash pain,<sup>63</sup> and work-related upper limb pain.<sup>64</sup> While the evidence favoring behavioral graded activity and cognition-targeted exercise therapy for patients with chronic pain was not generated from different cultural settings, the intervention is considered suitable for cultural adaptation. Future research should examine whether they are generic across cultures.

During cognition-targeted exercise therapy and graded exposure, the therapist continuously assesses, validates, and challenges the patients' culturally sensitive beliefs about pain and the anticipated outcome of each activity/exercise, to change maladaptive beliefs into positive ones. If certain activities/exercises are too threatening to perform, a preparatory phase of motor imagery can be included (including a home exercise program of motor imagery of the threatening exercise/activity).<sup>65</sup> Motor imagery is defined as a dynamic mental process involving the internal representation of an action without its actual motor output.<sup>66</sup> Although motor imagery seems to be a promising intervention that could be adapted in different cultural settings, consistent evidence from rigorous studies is still lacking.<sup>67</sup> The exercise/activity program should progress towards more feared movements and activities by discussing patients' fears and maladaptive perceptions, preferentially by asking questions that challenge the patients' (pain) beliefs. This part of the treatment is highly culturally-sensitive and allows the therapist to adapt to the patient's socio-cultural situation and interpretation of bodily responses to exercises and activities, including pain flares following the threatening exercises/activities. Therapists should try to decrease the anticipated danger (threat level) of the activities/exercises by challenging the nature of, and reasoning behind the patients' fears, assuring the safety of the activities/exercises, and increasing confidence in a successful accomplishment of the activities/exercise.<sup>50</sup> More details regarding this approach, including specific instructions on required communication skills, are described in detail elsewhere.<sup>50</sup>

## Challenges and strategies for cross-cultural pain education and exercises

Among the many challenges facing healthcare systems, is the major challenge of ensuring effective culturally congruent care (i.e., cultural diversity, sex, race, ethnicity, literacy) and linguistically appropriate information eliminating the disparities experienced by minority groups.<sup>68</sup> If cultural factors are not taken into account, the provision of care may be ineffective, if not harmful. Within each cultural group the pain experience includes pain expression, pain language, cultural context of suffering, traditional healers and lay remedies for pain, social roles/expectations, and perceptions of the health care system. Therefore, efforts should be made to produce and spread culturally adapted evidence-based materials and resources including language, content, and images that would resonate with the target population.<sup>69–72</sup>

Communication to the public aiming to improve knowledge and modify beliefs and attitudes about pain in multi-country public awareness campaigns (e.g., sponsored by Organizations) seems logical. However, the advantage associated with uniformity of practices and messages across countries is ambiguous, as it shows consistent global messages but limited adaptation to local situations.

During clinical practice, providing culturally appropriate services is, in many ways, more difficult because it may require clinicians adapting therapeutic interventions to

their culturally diverse patients.<sup>73</sup> There are still some concerns if cultural adaptation models that typically aim to adapt an intervention for one or more diverse groups in a way that retains fidelity to the core components or the original intervention should be used.<sup>72,74</sup> For example, researchers in the area of low back pain have identified 30 key messages that are important to deliver to patients with low back pain based on experts' and patients' perspectives.<sup>75</sup> However, it is unclear which of these target key messages produce clinically meaningful changes in pain and physical function or produce behavioral modifications to generate these changes.

Clinicians should be aware of patients' needs and goals and how cultural factors can shape patients' perceptions, conceptions, and unhelpful beliefs about exercises (e.g., fear of injury, negative consequences). For instance, adherence to exercises can be influenced by firm rules and norms about gender roles in some cultures (e.g., women are responsible for household duties, exercise is regarded as a 'masculine' and inappropriate practice for women)<sup>76</sup> and also by psychological barriers such as depression (e.g., post-war trauma).<sup>77</sup> Clinicians may have difficulties in challenging unhelpful beliefs, enhancing self-efficacy, providing safety-cues, providing advice on suitable levels of pain, and providing advice on exercise modification where language difference is an issue. In these cases, a peer-led exercise and education program can be effectively utilized to overcome language and cultural barriers, to alleviate pressurized health care professionals, and to facilitate behavior change.<sup>56,78</sup> Another aspect to be considered is that some patients require privacy to do exercises, while others prefer to engage in group exercise sessions composed of individuals of the same community where they feel safe and accepted.<sup>79</sup> Strategies for the development of culturally adapted educational and exercise interventions are presented in the [Supplementary Material](#).

## Future directions

Cross-cultural adaptations (i.e., linguistically appropriate and culturally sensitive) are necessary before using these interventions in a population which is different from the one used to develop the intervention. Clinicians should be trained to deliver a culturally sensitive approach in their practice and programs developed for minority groups might help to decrease health inequalities. Currently, few studies have investigated the feasibility<sup>36,80</sup> and effectiveness of a culturally-sensitive pain education intervention in patients with (persistent) pain.<sup>72</sup> Further research, especially including non-Western populations and minority groups (e.g., migrants and refugees), are essential to determine the extent to which pain treatments may need to be culturally adapted to make them most appropriate to new populations who may live in different countries or speak different languages than those for whom the interventions were first developed. In addition to the growing concerns about evidence-based interventions in the pain field, attention to cultural diversity is crucial and should start in the entry-level physical therapy program. It would be valuable to integrate students and community members from a variety of

ethnocultural backgrounds, embracing the diversity in the classroom to provide opportunities for experiential learning.<sup>12</sup>

## Conclusion

Maladaptive pain beliefs and behaviors can be shaped by patients' culture. Clinicians should consider patients' beliefs, values, and practices to provide individualized care to their culturally diverse patients. In this aspect, the contents of pain education should be presented using different culturally appropriate examples, metaphors, images, and delivery methods that may enhance the impact of the message. A culturally sensitive approach may help to introduce patients to graded activities, so that they can apply these strategies in culturally acceptable and meaningful ways. Future studies should investigate the effectiveness of culturally-adapted interventions in pain-related outcomes in different pain conditions in patients with different cultural backgrounds.

## Conflicts of interest

The Vrije Universiteit Brussel (VUB) and Jo Nijs received lecturing fees.

Jo Nijs co-authored a Dutch book on pain neuroscience education; the royalties are collected by the VUB.

## Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.bjpt.2022.100442](https://doi.org/10.1016/j.bjpt.2022.100442).

## References

- Vos T, Allen C, Arora M, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the global burden of disease study 2015. *Lancet*. 2016;388(10053):1545–1602. [https://doi.org/10.1016/S0140-6736\(16\)31678-6](https://doi.org/10.1016/S0140-6736(16)31678-6).
- Raja SN, Carr DB, Cohen M, et al. The revised international association for the study of pain definition of pain: concepts, challenges, and compromises. *Pain*. 2020;161(9):1976–1982.
- Gatchel RJ, Peng YB, Peters ML, Fuchs PN, Turk DC. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychol Bull*. 2007;133(4):581. <https://doi.org/10.1037/0033-2909.133.4.581>.
- Wijma AJ, van Wilgen CP, Meeus M, Nijs J. Clinical biopsychosocial physiotherapy assessment of patients with chronic pain: the first step in pain neuroscience education. *Physiother Theory Pract*. 2016;32(5):368–384.
- Karran EL, Grant AR, Moseley GL. Low back pain and the social determinants of health: a systematic review and narrative synthesis. *Pain*. 2020;161(11):2476–2493.
- Stilwell P, Harman K. An enactive approach to pain: beyond the biopsychosocial model. *Phenomenol Cogn Sci*. 2019;18(4):637–665.
- Mescouto K, Olson RE, Hodges PW, Setchell J. A critical review of the biopsychosocial model of low back pain care: time for a new approach? *Disabil Rehabil*. 2020;7:1–15.
- Mardian AS, Hanson ER, Villarroel L, et al. Flipping the pain care model: a sociopsychobiological approach to high-value chronic pain care. *Pain Med*. 2020;21(6):1168–1180.
- Meints SM, Miller MM, Hirsh AT. Differences in pain coping between black and white Americans: a meta-analysis. *J Pain*. 2016;17(6):642–653.
- Orhan C, Van Looveren E, Cagnie B, Mukhtar NB, Lenoir DMM. Are pain beliefs, cognitions, and behaviors influenced by race, ethnicity, and culture in patients with chronic musculoskeletal pain: a systematic review. *Pain Physician*. 2018;21(6):541–558.
- UNESCO. The 2009 UNESCO framework for cultural statistics (FCS). Published 2009. Accessed 6 January 2021. <https://unstats.un.org/unsd/statcom/doc10/BG-FCS-E.pdf>.
- Brady B, Veljanova I, Chipchase LS. Culturally informed practice and physiotherapy. *J Physiother*. 2016;62(3):121–123.
- Caneiro JP, Bunzli S, O'Sullivan P. Beliefs about the body and pain: the critical role in musculoskeletal pain management. *Braz J Phys Ther*. 2020;25(1):17–29.
- Leeuw M, Goossens MEJB, Linton SJ, Crombez G, Boersma K, Vlaeyen JWS. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med*. 2007. <https://doi.org/10.1007/s10865-006-9085-0>. Published online.
- Quartana PJ, Campbell CM, Edwards RR. Pain catastrophizing: a critical review. *Expert Rev Neurother*. 2009;9(5):745–758.
- Vlaeyen JWS, Crombez G, Linton SJ. The fear-avoidance model of pain. *Pain*. 2016;157(8):1588–1589. <https://doi.org/10.1097/j.pain.0000000000000574>.
- Bailey R, Wells A. Metacognitive beliefs moderate the relationship between catastrophic misinterpretation and health anxiety. *J Anxiety Disord*. 2015;34:8–14.
- Büssing A, Ostermann T, Neugebauer EAM, Heusser P. Adaptive coping strategies in patients with chronic pain conditions and their interpretation of disease. *BMC Public Health*. 2010;10(1):1–10.
- Sharma S, Ferreira-Valente A, de C Williams AC, Abbott JH, Pais-Ribeiro J, Jensen MP. Group differences between countries and between languages in pain-related beliefs, coping, and catastrophizing in chronic pain: a systematic review. *Pain Med*. 2020;21(9):1847–1862.
- Lin I, Wiles L, Waller R, et al. What does best practice care for musculoskeletal pain look like? Eleven consistent recommendations from high-quality clinical practice guidelines: systematic review. *Br J Sports Med*. 2020;54(2):79–86. <https://doi.org/10.1136/bjsports-2018-099878>.
- Brady B, Veljanova I, Chipchase L. Are multidisciplinary interventions multicultural? A topical review of the pain literature as it relates to culturally diverse patient groups. *Pain*. 2016;157(2):321–328.
- Edwards CL, Fillingim RB, Keefe F. Race, ethnicity and pain. *Pain*. 2001;94(2):133–137.
- Kim HJ, Yang GS, Greenspan JD, et al. Racial and ethnic differences in experimental pain sensitivity: systematic review and meta-analysis. *Pain*. 2017;158(2):194–211.
- Boerner KE, Chambers CT, Gahagan J, Keogh E, Fillingim RB, Mogil JS. Conceptual complexity of gender and its relevance to pain. *Pain*. 2018;159(11):2137–2141.
- Green CR, KO Anderson, Baker TA, et al. The unequal burden of pain: confronting racial and ethnic disparities in pain. *Pain Med*. 2003;4(3):277–294.
- Myers CD, Riley III JL, Robinson ME. Psychosocial contributions to sex-correlated differences in pain. *Clin J Pain*. 2003;19(4):225–232.
- Anderson SR, Losin EAR. A sociocultural neuroscience approach to pain. *Cult Brain*. 2017;5(1):14–35.
- Honeyman PT, Jacobs EA. Effects of culture on back pain in Australian aboriginals. *Spine (Phila Pa 1976)*. 1996;21(7):841–843.

29. Madden V, O'Sullivan P, Fisher J, Malambule B. Our training left us unprepared—two physiotherapists' reflections after working with women with low back pain in a rural Zulu community in South Africa. *J Community Heal Sci*. 2013;8(2).
30. Ferreira-Valente A, Sharma S, Torres S, et al. Does religiosity/spirituality play a role in function, pain-related beliefs, and coping in patients with chronic pain? A systematic review. *J Relig Health*. 2019;1–55. Published online.
31. Juarez G, Ferrell B, Borneman T. Influence of culture on cancer pain management in Hispanic patients. *Cancer Pract*. 1998;6(5):262–269.
32. Arraras JI, Wright SJ, Jusue G, Tejedor M, Calvo JI. Coping style, locus of control, psychological distress and pain-related behaviours in cancer and other diseases. *Psychol Health Med*. 2002;7(2):181–187.
33. Lin IB, O'Sullivan PB, Coffin JA, Mak DB, Toussaint S, Straker LM. Disabling chronic low back pain as an iatrogenic disorder: a qualitative study in Aboriginal Australians. *BMJ Open*. 2013;3(4): e002654.
34. Sharma S, Abbott JH, Jensen MP. Why clinicians should consider the role of culture in chronic pain. *Braz J Phys Ther*. 2018;22(5):345.
35. Anderson RT. An orthopedic ethnography in rural Nepal. *Med Anthropol*. 1984;8(1):46–59.
36. Sharma S, Jensen MP, Moseley GL, Abbott JH. Results of a feasibility randomised clinical trial on pain education for low back pain in Nepal: the Pain Education in Nepal-Low Back Pain (PEN-LBP) feasibility trial. *BMJ Open*. 2019;9(3): e026874.
37. Smith JA, Osborn M. Pain as an assault on the self: an interpretative phenomenological analysis of the psychological impact of chronic benign low back pain. *Psychol Heal*. 2007;22(5):517–534.
38. Fabian LA, McGuire L, Goodin BR, Edwards RR. Ethnicity, catastrophizing, and qualities of the pain experience. *Pain Med*. 2011;12(2):314–321.
39. Forsythe LP, Thorn B, Day M, Shelby G. Race and sex differences in primary appraisals, catastrophizing, and experimental pain outcomes. *J pain*. 2011;12(5):563–572.
40. Shaw WS, Campbell P, Nelson CC, Main CJ, Linton SJ. Effects of workplace, family and cultural influences on low back pain: what opportunities exist to address social factors in general consultations? *Best Pract Res Clin Rheumatol*. 2013;27(5):637–648.
41. Lehman BJ, David DM, Gruber JA. Rethinking the biopsychosocial model of health: understanding health as a dynamic system. *Soc Personal Psychol Compass*. 2017;11(8):e12328.
42. De Ruddere L, Craig KD. Understanding stigma and chronic pain: a state-of-the-art review. *Pain*. 2016;157(8):1607–1610.
43. O'Keefe M, O'Sullivan P, Purtill H, Bargary N, O'Sullivan K. Cognitive functional therapy compared with a group-based exercise and education intervention for chronic low back pain: a multicentre randomised controlled trial (RCT). *Br J Sports Med*. 2020;54(13):782–789.
44. Castro J, Correia L, Donato BS, et al. Cognitive functional therapy compared with core exercise and manual therapy in patients with chronic low back pain: randomised controlled trial. *Pain*. 2022. <https://doi.org/10.1097/j.pain.0000000000002644>. Published online.
45. Lamb SE, Hansen Z, Lall R, et al. Group cognitive behavioural treatment for low-back pain in primary care: a randomised controlled trial and cost-effectiveness analysis. *Lancet*. 2010;375(9718):916–923. [https://doi.org/10.1016/S0140-6736\(09\)62164-4](https://doi.org/10.1016/S0140-6736(09)62164-4).
46. de C Williams AC, Fisher E, Hearn L, Eccleston C. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database Syst Rev*. 2020(8).
47. Soares LO, Ferreira GE, Costa LOP, Nogueira LC, Meziat-Filho N, Reis FJJ. Meditation for adults with non-specific low back pain: a systematic review and meta-analysis. *Scand J Pain*. 2022;22(1):26–39.
48. Schütze R, Rees C, Smith A, Slater H, Campbell JM, O'Sullivan P. How can we best reduce pain catastrophizing in adults with chronic noncancer pain? A systematic review and meta-analysis. *J Pain*. 2018;19(3):233–256.
49. Wood L, Hendrick PA. A systematic review and meta-analysis of pain neuroscience education for chronic low back pain: short- and long-term outcomes of pain and disability. *Eur J Pain*. 2019;23(2):234–249. <https://doi.org/10.1002/ejp.1314>.
50. Nijs J, Lluch Gires E, Lundberg M, Malfliet A, Sterling M. Exercise therapy for chronic musculoskeletal pain: innovation by altering pain memories. *Man Ther*. 2015;20(1):216–220. <https://doi.org/10.1016/j.math.2014.07.004>.
51. Macedo LG, Latimer J, Maher CG, et al. Motor control or graded activity exercises for chronic low back pain? A randomised controlled trial. *BMC Musculoskelet Disord*. 2008;9(1):65.
52. Vlaeyen JWS, de Jong J, Geilen M, Heuts PHTG, van Breukelen G. The treatment of fear of movement/(re) injury in chronic low back pain: further evidence on the effectiveness of exposure in vivo. *Clin J Pain*. 2002;18(4):251–261.
53. Watson JA, Ryan CG, Cooper L, et al. Pain neuroscience education for adults with chronic musculoskeletal pain: a mixed-methods systematic review and meta-analysis. *J Pain*. 2019;20(10):1140.e1–1140.e22. <https://doi.org/10.1016/j.jpain.2019.02.011>.
54. Wood L, Hendrick PA. A systematic review and meta-analysis of pain neuroscience education for chronic low back pain: short- and long-term outcomes of pain and disability. *Eur J Pain*. 2018;23(2):234–249. <https://doi.org/10.1002/ejp.1314>.
55. Lim YZ, Chou L, Au RTM, et al. People with low back pain want clear, consistent and personalised information on prognosis, treatment options and self-management strategies: a systematic review. *J Physiother*. 2019;65(3):124–135.
56. Parker R, Aidsma J, Stein DJ. Managing pain in women living with HIV/AIDS: a randomized controlled trial testing the effect of a six-week peer-led exercise and education intervention. *J Nerv Ment Dis*. 2016;204(9):665–672.
57. Malfliet A, Kregel J, Coppieters I, et al. Effect of pain neuroscience education combined with cognition-targeted motor control training on chronic spinal pain: a randomized clinical trial. *JAMA Neurol*. 2018;Jul;17(7):808–817.
58. Nijs J, D'Hondt E, Clarys P, et al. Lifestyle and chronic pain across the lifespan: an inconvenient truth? *PMR*. 2020;12(4):410–419.
59. Veenhof C, Köke AJA, Dekker J, et al. Effectiveness of behavioral graded activity in patients with osteoarthritis of the hip and/or knee: a randomized clinical trial. *Arthritis Care Res Off J Am Coll Rheumatol*. 2006;55(6):925–934.
60. López-de-Uralde-Villanueva I, Muñoz-García D, Gil-Martínez A, et al. A systematic review and meta-analysis on the effectiveness of graded activity and graded exposure for chronic nonspecific low back pain. *Pain Med*. 2016;17(1):172–188.
61. Leeuw M, Goossens MEJB, van Breukelen GJP, et al. Exposure in vivo versus operant graded activity in chronic low back pain patients: results of a randomized controlled trial. *Pain*. 2008;138(1):192–207.
62. de Jong JR, Vlaeyen JWS, Onghena P, Cuypers C, den Hollander M, Ruijgrok J. Reduction of pain-related fear in complex regional pain syndrome type I: the application of graded exposure in vivo. *Pain*. 2005;116(3):264–275.
63. de Jong JR, Vangronsveld K, Peters ML, et al. Reduction of pain-related fear and disability in post-traumatic neck pain: a replicated single-case experimental study of exposure in vivo. *J Pain*. 2008;9(12):1123–1134.
64. de Jong JR, Vlaeyen JWS, van Eijsden M, Loo C, Onghena P. Reduction of pain-related fear and increased function and participation in work-related upper extremity pain (WRUEP): effects of exposure in vivo. *PAIN®*. 2012;153(10):2109–2118.
65. Sengul YS, Kaya N, Yalcinkaya G, Kirmizi M, Kalemci O. The effects of the addition of motor imagery to home exercises on pain, disability and psychosocial parameters in patients



- undergoing lumbar spinal surgery: a randomized controlled trial. *EXPLORE*. 2021;17(4):334–339.
66. Decety J. The neurophysiological basis of motor imagery. *Behav Brain Res*. 1996;77(1–2):45–52.
  67. Suso-Martí L, La Touche R, Angulo-Díaz-Parreño S, Cuenca-Martínez F. Effectiveness of motor imagery and action observation training on musculoskeletal pain intensity: a systematic review and meta-analysis. *Eur J Pain*. 2020;24(5):886–901.
  68. Castro FG, Barrera Jr M, Holleran Steiker LK. Issues and challenges in the design of culturally adapted evidence-based interventions. *Annu Rev Clin Psychol*. 2010;6:213–239.
  69. Reis FJJ, Bengaly AGC, Valentim JCP, et al. An E-Pain intervention to spread modern pain education in Brazil. *Braz J Phys Ther*. 2017;21(5):305–306. <https://doi.org/10.1016/j.bjpt.2017.06.020>.
  70. Reis F, Palermo TM, Acalantis L, et al. “A journey to learn about pain”: the development and validation of a comic book about pain neuroscience education for children. *Brazilian J Phys Ther*. 2021;26(1): 100348. <https://doi.org/10.1016/j.bjpt.2021.04.009>.
  71. Pas R, Meeus M, Malfliet A, et al. Development and feasibility testing of a pain neuroscience education program for children with chronic pain: treatment protocol. *Braz J Phys Ther*. 2018;22(3):248–253. <https://doi.org/10.1016/j.bjpt.2018.02.004>.
  72. Orhan C, Cagnie B, Favoreel A, et al. Development of culturally sensitive pain neuroscience education for first-generation Turkish patients with chronic pain: a modified Delphi study. *Musculoskelet Sci Pract*. 2019;39:1–9.
  73. Narayan MC. Six steps towards cultural competence: a clinician’s guide. *Home Health Care Manag Pract*. 2002;14(5):378–386.
  74. Gonzales NA. Expanding the cultural adaptation framework for population-level impact. *Prev Sci*. 2017;18(6):689–693.
  75. French SD, Nielsen M, Hall L, et al. Essential key messages about diagnosis, imaging, and self-care for people with low back pain: a modified Delphi study of consumer and expert opinions. *Pain*. 2019;160(12):2787–2797.
  76. Edelstein OE, Vered I, Sarid O. Correlates of participation in physical activity among older women in Israel: does ethno-cultural background matter? *Health Promot Int*. 2021;36(1):34–45.
  77. Caperchione CM, Kolt GS, Tennent R, Mummery WK. Physical activity behaviours of culturally and linguistically diverse (CALD) women living in Australia: a qualitative study of socio-cultural influences. *BMC Public Health*. 2011;11(1):26.
  78. Saw MM, Kruger-Jakins T, Edries N, Parker R. Significant improvements in pain after a six-week physiotherapist-led exercise and education intervention, in patients with osteoarthritis awaiting arthroplasty, in South Africa: a randomised controlled trial. *BMC Musculoskelet Disord*. 2016;17(1):236.
  79. Medagama A, Galgomuwa M. Lack of infrastructure, social and cultural factors limit physical activity among patients with type 2 diabetes in rural Sri Lanka, a qualitative study. *PLoS ONE*. 2018;13(2): e0192679.
  80. Mukhtar NB, Meeus M, Gursen C, Mohammed J, De Pauw R, Cagnie B. Pilot study on the effects of a culturally-sensitive and standard pain neuroscience education for Hausa-speaking patients with chronic neck pain. *Disabil Rehabil*. 2021:1–11. Published online.