

Methods: This is a cross-sectional analytical study. The study included women using hormone therapy (Tamoxifen or Anastrozole) for breast cancer and complaining of arthralgia (at least 1 point on the Numerical Categorical Scale - NCS and 1 joint located on the Brief Pain Inventory - BPI). Women who had stage IV cancer, with the presence of lymphedema, limitation to answer questionnaires and women with tumor recurrence were excluded. Data distribution was evaluated using the Kolmogorov-Smirnov test. To determine the isolated contribution of Hormone Therapy Time (independent variable) to arthralgia (dependent variable) in women of hormone therapy for breast cancer, linear regression analyzes, and Spearman correlation coefficients were performed. The significance level was set at 5%. The correlation was classified according to the following criteria: weak (0.0 - 0.4), moderate (0.4 - 0.7) and strong (0.7 - 1.0). **Results:** ninety-two women with a mean age of 53.68 years and a standard deviation of 9.53 participated in the study. The mean duration of hormone therapy use was 2.44 years, with a standard deviation of 1.45, and the mean duration of arthralgia was 5.38. with a standard deviation of 2.73. Arthralgia and duration of hormone therapy did not present a significant correlation ($p=0.11$; $r=0.16$). **Conclusion:** no association was observed between the duration of hormone therapy and arthralgia in women undergoing treatment for breast cancer.

Implications: Although numerous studies report the presence of arthralgia in women who use hormone therapy in the treatment of breast cancer, the duration of hormone therapy does not seem to influence this complaint. However, prospective cohort studies are needed to confirm the effects of hormone therapy duration on arthralgia in women undergoing cancer treatment.

Keywords: Arthralgia, Hormone therapy, Association

Conflict of interest: The authors declare no conflict of interest.

Acknowledgment: Not applicable.

Ethics committee approval: Faculty of Ceiland, University of Brasília - UnB, approval opinion n° 3.022.045.

<https://doi.org/10.1016/j.bjpt.2024.100675>

79

FLEXIBILITY AFTER 16 WEEKS OF PILATES EXERCISE IN WOMEN WITH ARTHRALGIA AFTER HORMONE THERAPY FOR BREAST CANCER

Liana Matheus¹, Cariele Freitas¹, Keyla Barbosa¹, Lidiane da Silva², Patrícia Garcia¹, Simone Botelho³

¹ Department of Physical Therapy, University of Brasília (UnB), Brasília, Distrito Federal, Brazil

² University Hospital of Brasília, Brasília, Distrito Federal, Brazil

³ Department of Physical Therapy, Federal University of Alfenas (UNIFAL), Alfenas, Minas Gerais, Brazil

Background: arthralgia is a common symptom after treatment with hormone therapy for breast cancer and can lead to several complications, including loss of flexibility. Studies have recommended performing the Pilates exercise to minimize morbidities resulting from cancer treatment.

Objectives: to evaluate flexibility after 16 weeks of Pilates exercise in women with arthralgia after hormone therapy for breast cancer.

Methods: observational study, carried out at the physiotherapy clinic of the High Complexity Oncology Unit. Women with complaints of arthralgia during hormone therapy and who participated in a Mat Pilates exercise program (twice a week for 16 weeks) were included, and women with active cancer or who did not complete the 16 weeks of exercise were excluded. Flexibility was evaluated through the "sit-and-reach test" using the SANNY Instant Unisex Pro

Portable Wells Bench. Statistical analysis was performed using GraphPad Prism. The data were submitted to the Kolmogorov-Smirnov normality test, followed by the paired t-test (parametric data) or the Wilcoxon test for repeated measures (non-parametric data), considering a significance level of 5%.

Results: eight women with a mean age of 57.25 ± 12.61 completed the 16 weeks of performing the Pilates exercise, 5 using hormone therapy with Tamoxifen and 3 using Anastrozole. The women had mean and standard deviation for flexibility of 18.12 ± 5.66 at the beginning and 22.75 ± 6.88 after 16 weeks of Pilates exercise ($p=0.004$). The group that used Tamoxifen presented flexibility of 16.7 ± 4.99 before and 21.4 ± 7.57 after Pilates ($p=0.04$) and the group using Anastrozole presented flexibility of 20.5 ± 7 before and 25 ± 6.26 after Pilates ($p=0.16$).

Conclusion: There was greater flexibility in women after performing the Pilates exercise.

Implications: Pilates exercises have the potential to improve the flexibility of women undergoing cancer treatment. However, randomized controlled trials, with a significant sample, are needed for better scientific evidence of the effects of Pilates exercise on the flexibility of women with arthralgia.

Keywords: Arthralgia, Pilates, Pliability

Conflict of interest: The authors declare no conflict of interest.

Acknowledgment: Not applicable.

Ethics committee approval: University of Brasília, n° 3.022.045.

<https://doi.org/10.1016/j.bjpt.2024.100676>

80

ELECTROMYOGRAPHIC PROFILE OF THE WRIST AND ELBOW FLEXORS DURING PNF MOTOR IRRADIATION

Carlos Henrique R Horsczaruk¹, José Vicente P Martins², Claudia Vargas², Laura A.S Oliveira¹, Thiago C Lemos¹

¹ Programa de Pós-Graduação em Ciências da Reabilitação, Centro Universitário Augusto Motta (UNISUAM), Rio de Janeiro, Rio de Janeiro, Brasil

² Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Rio de Janeiro, Brasil

Background: Proprioceptive Neuromuscular Facilitation (PNF) is a concept that has been applied to treating numerous disabling disorders. Among the basic PNF procedures, there is motor irradiation, in which resistance is applied to a body segment to generate muscle activation in another segment and thus obtain improvement in muscle strength. The generated muscular activation can be analyzed in several ways, one of them being Electromyography (EMG). Electromyographic analysis has suggested that target muscles of irradiation demonstrate electromyographic activity during irradiation in healthy individuals and in those with neurological disorders. Despite this, the neuromuscular activation profile (amplitude and temporality) resulting from the different stimuli and positions used in PNF has not yet been investigated.

Objectives: To analyze the neuromuscular activation profile of upper limb muscles in healthy individuals during the application of different PNF irradiation protocols.

Methods: This is a cross-sectional study in which 32 healthy individuals of both sexes, aged between 18 and 45 years. After signing the consent form, the handedness of a subject in activities of daily living, level of physical activity, and sociodemographic characteristics will be identified. First, the EMG signals of the upper limb muscles (biceps brachii, brachioradialis, flexor carpi radialis, and triceps) will be collected during a maximum voluntary contraction (3 repetitions held for 3s with an interval of 30s between contractions).