

Original Research

Five-repetition chair-stand test vs. handgrip strength: Which better predicts mortality risk? a follow-up study in 43,605 middle-aged and older adults

Rodrigo Núñez-Cortés^a, Lars Louis Andersen^b, Carlos Cruz-Montecinos^{a,*}, Ana Polo-López^c, Rubén López-Bueno^{b,c,d}, Joaquín Calatayud^{b,d}^a Department of Physical Therapy, Faculty of Medicine, University of Chile, Santiago, Chile^b National Research Centre for the Working Environment, Copenhagen, Denmark^c Exercise Intervention for Health Research Group (EXINH-RG), Department of Physiotherapy, University of Valencia, Spain^d Department of Physical Medicine and Nursing, University of Zaragoza, Zaragoza, Spain

ARTICLE INFO

Keywords:

Aging population

Functional performance

Physical fitness markers

Sarcopenia

Survival Analysis

ABSTRACT

Background: Ageing reduces muscle strength and function, increasing mortality risk. Identifying simple performance markers can guide interventions for healthy ageing.**Objective:** To assess the prospective dose-response association of the 5-repetition Chair Stand Test (5-CST) and handgrip strength (HGS) with mortality in middle-aged and older adults.**Methods:** This prospective study included community-dwelling participants aged 50 years or older from the SHARE study. HGS and 5-CST were assessed at baseline, with all-cause mortality tracked through follow-up interviews. Cox regression with restricted cubic splines was used, controlling for several confounders.**Results:** 43,605 participants (mean age (SD): 65.3 (9.1), 54 % women) were included. During a mean follow-up of 7.3 ± 2.2 years, there were 4154 deaths (9.5 %). Both 5-CST and HGS were curvilinearly associated with all-cause mortality. Using the median level of 5-CST as a reference (11 s), 10th percentile of 5-CST (7 s) showed a hazard ratio (HR) of 0.74 (95 %CI: 0.69, 0.80). The 90th percentile (18 s) of 5-CST showed a HR of 1.18 (95 %CI: 1.14, 1.22). Stratified analysis indicated 5-CST was most strongly associated with mortality in women. Regarding HGS, using the median level as a reference (33 kg), the 10th percentile of muscle strength (21 kg) showed a HR of 1.62 (95 %CI: 1.50, 1.75). The 90th percentile (51 kg) of muscle strength showed a HR of 0.58 (95 %CI: 0.52, 0.64).**Conclusion:** Both tools provide valuable information, but HGS may be considered more relevant for identifying those at increased mortality risk, while 5-CST may be especially useful in women.

Introduction

Aging is associated with physiological and architectural changes in musculoskeletal health,¹ including decreased muscle mass and increased intramuscular fat.² This decline can have a significant impact on daily activities.³ In this regard, rapid population aging is a global phenomenon that poses important public health challenges,⁴ such as the need to identify markers to predict and prevent mortality risk and guide the design of interventions to promote healthy aging. The implementation of preventive strategies aimed at improving physical health can not only reduce mortality in older adults but also enhance quality of life. Among physical health indicators, low levels of muscle strength and functional capacity are considered important risk factors for morbidity and

mortality in this age group.

Handgrip strength (HGS) and the 5-repetition Chair Stand Test (5-CST) are two widely used tests to assess physical health in older people.^{5,6} HGS is an inexpensive, rapid, simple, and reliable marker of general muscular strength in adults.⁷⁻⁹ Moreover, HGS is widely accepted as an explicator of frailty, cognitive impairment, multimorbidity, and all-cause and disease-specific mortality.¹⁰⁻¹³ On the other hand, the 5-CST assesses functional capacity, fall risk, and lower-limb strength, providing a more integrated view of mobility and functionality.¹⁴⁻¹⁶ Importantly, a major advantage of 5-CST is that it is easy to perform in a variety of settings, including the home, as no other equipment than a timer and a chair is required. Both tests are endorsed by major clinical guidelines and research frameworks, such as the

* Corresponding author at: Department of Physical Therapy, Faculty of Medicine, University of Chile, Av. Independencia 1027, Santiago, Chile.

E-mail address: carloscruz@uchile.cl (C. Cruz-Montecinos).<https://doi.org/10.1016/j.bjpt.2025.101529>

Received 15 February 2025; Received in revised form 23 July 2025; Accepted 14 August 2025

1413-3555/© 2025 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier España, S.L.U. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

EWGSOP2 consensus,¹⁷ as key components in the assessment of sarcopenia, frailty, and functional decline. Their widespread use in epidemiologic studies facilitates international comparisons and benchmarking. In addition, both the HGS and the 5-CST have shown consistent associations with adverse outcomes such as disability, hospitalization, and mortality, making them particularly valuable for public health surveillance and prognostic modeling. Nevertheless, few studies have directly compared the predictive value of both tests within the same cohort, leaving an important gap in understanding their relative utility for mortality risk stratification.

Given their simplicity and applicability in clinical and research settings, the debate over which of these two tests best predicts mortality remains a relevant topic.^{18,19} Mainly, for the use of HGS as an indicator of global strength, considering the conflicting evidence of association between HGS with lower limb strength (e.g., knee extensor) regardless of age and health status.¹⁸ Thus, evaluating the prospective association of HGS and 5-CST with mortality in the elderly may provide a basis for optimizing assessment tools and guiding interventions to prevent functional decline and reduce mortality risk. This study aims to compare the predictive dose-response ability of HGS and 5-CST with respect to mortality in a large, multi-country representative sample of middle-aged and older adults. The value of each test separately in its prognostic value for mortality is expected to be known.

Methods

Participants and design

This prospective study uses data from the Survey of Health, Ageing and Retirement in Europe (SHARE), covering 27 European countries and Israel.^{20,21} Among all observations in the dataset ($n = 685,919$), we included a total of 43,605 community-dwelling individuals aged ≥ 50 years who had valid measurements of both HGS and 5-CST at wave 5 (baseline, 2013), along with data from at least one follow-up assessment up to December 2022. A detailed flowchart of the selection process is shown in Fig. 1.

SHARE employs a multi-stage stratified sampling design in which participating countries are divided into different strata according to geographical area, and municipalities or postcodes within these strata serve as primary sampling units.^{20,21} Data are collected in each survey wave through computer-assisted face-to-face interviews in the home. SHARE uses pre-harmonized interviews, and new respondents are enrolled in each wave to compensate for losses. The target population of SHARE is everyone aged 50 and over whose usual place of residence is in

one of the countries participating in SHARE. Individuals are excluded from the initial or refreshment sample if they are imprisoned, hospitalized, or out of the country during the entire survey period, if they do not speak the language(s) of the country, or if they have moved to an unknown address. Regardless of their age, partners living in the same household are interviewed.

Chair stand test

The 5-CST was measured once per participant. Participants were instructed to cross their arms over their chest, stand up from a seated position in a chair, and then sit down again, repeating this sequence five times as quickly as possible. A stopwatch was used to measure the time (in seconds) taken to complete five repetitions without using the hands.

Handgrip strength

HGS was assessed using a handgrip dynamometer (Smedley, S Dynamometer, TTM, Tokyo, 0–100 kg). Each hand was measured twice under the guidance of trained interviewers. Participants held their upper arm parallel to the trunk, with the elbow at 90° flexion and the wrist in a neutral position while standing or sitting. Trained interviewers provided standardized instructions to press the dynamometer with maximum effort for 2 s. The value of HGS (unit: kg) was defined as the maximum value of either hand for the present analyses.

Mortality

Death was recorded in end-of-life interviews with proxy respondents, that is, a relative, household member, neighbor, or other person close to the deceased participant, who provided information on date and cause of death. For cause of death, the proxy respondent was asked "What was the main cause of death?" with multiple response options. For this study, we focused on all-cause mortality, which includes deaths from any cause. In the case of a missing value for the date of death, we imputed the mean of the date of the last interview with the participant and the date of the interview with the end-of-life representative. This imputation was applied in 3.3 % of all recorded deaths.

Covariates

The potential confounders included in the adjustment were: age, sex, body mass index (BMI), smoking status (current smoker, former smoker, never smoker), educational level (classified as low, medium, or high based on ISCED-1997 categories), marital status (divorced or married, living separated from spouse, never married or widowed), geographic region (Eastern, Northern, Southern, Western Europe, and Israel), and number of medications taken.

Statistical analyses

Cox proportional hazards regression was used to assess the prospective association of 5-CST and HGS, respectively, with all-cause mortality. We used restricted cubic splines with knots at the 10th, 50th, and 90th percentiles of the 5-CST time and HGS kilogram distribution to account for potential nonlinearity. Results were presented as hazard ratios (HR) with 95 % confidence intervals (CI). All analyses were adjusted for age, sex, body mass index, smoking status, educational level, marital status, geographic region, and number of medications taken.

To visualize the association of 5-CST and HGS with the risk of all-cause mortality, we plotted the estimated HRs against the predictors, using the median values as the reference point. In addition, a stratified analysis by sex was performed. All statistical analyses were performed with SAS version 9.4 (SAS Institute Inc., Cary, NC, USA). A p-value of less than 0.05 was considered statistically significant.




Fig. 1. Flowchart of Participant Selection. Abbreviations: Handgrip Strength (HGS) and Five Repetition Chair Stand Test (5-CST).

Results

The study included 43,605 participants with a mean age of 65.3 years (SD = 9.1). The sample was 54 % female and 46 % male. The mean BMI was 26.7 kg/m² (SD 4.5). The cohort was geographically diverse, with 44.4 % from Western Europe, 23.8 % from Northern Europe, 20.8 % from Southern Europe, 8.0 % from Eastern Europe, and 3.0 % from Israel. The baseline characteristics of the participants are detailed in Table 1. During a mean follow-up of 7.3 ± 2.2 years, a total of 4154 deaths (9.5 %) from all causes were recorded.

A curvilinear association between performance on the 5-CST and the risk of all-cause mortality was observed. As illustrated in Fig. 2, using the median 5-CST time (11 s) as the reference point, individuals in the 10th percentile (7 s) had a hazard ratio (HR) of 0.74 (95 % CI: 0.69, 0.80), indicating a protective effect. In contrast, those in the 90th percentile (18 s) had a HR of 1.18 (95 % CI: 1.14, 1.22), reflecting increased risk.

Sex-stratified analyses revealed differential patterns in the 5-CST–mortality relationship. As shown in Fig. 3, while both men and women demonstrated increased risk with slower performance, the association was more pronounced in women. In men, the relationship plateaued beyond 17 s (90th percentile = 17 s, HR = 1.19, 95 % CI: 1.13, 1.24), whereas in women, the risk continued to rise (90th percentile = 19 s, HR = 1.24, 95 % CI: 1.17, 1.32).

Fig. 4 displays the relationship between handgrip strength (HGS) and all-cause mortality, which also followed a curvilinear pattern. Using the median grip strength (33 kg) as the reference, individuals in the 10th percentile (21 kg) exhibited a significantly elevated risk (HR = 1.62, 95 % CI: 1.50, 1.75), while those in the 90th percentile (51 kg) had a lower

Table 1
Baseline characteristics of participants (n = 43,605).

		n	%	Mean (SD)
Age (years)			65.3 (9.1)	
Body mass index (kg/m ²)			26.7 (4.5)	
Sex				
	Male	20,039	46.0	
	Female	23,566	54.0	
Smoking				
	Smoker	7775	17.8	
	Ex-smoker	13,506	31.0	
	Never smoked	22,324	51.2	
Education				
	Low education	15,189	34.8	
	Medium education	17,192	39.4	
	High education	11,224	25.7	
Marital status				
	Married and living together with spouse OR Registered partnership	31,890	73.1	
	Divorced OR Married, living separated from spouse	4448	10.2	
	Never married	2400	5.5	
	Widowed	4867	11.2	
Geographic Region (United Nations definition)				
	Eastern Europe	3474	8.0	
	Northern Europe	10,390	23.8	
	Southern Europe	9062	20.8	
	Western Europe	19,379	44.4	
	Israel	1300	3.0	
Number of drugs				
	0	12,155	27.9	
	1	12,814	29.4	
	2	8684	19.9	
	3	5239	12.0	
	4	2622	6.0	
	5	2091	4.8	




Fig. 2. Association of 5-repetition Chair Stand Test with risk of death during follow-up in adults older than 50 years (n = 43,605). Note: The solid line shows the hazard ratios and the dashed lines show the 95 % confidence intervals. Model adjusted for age, sex, body mass index, smoking status, educational level, marital status, geographic region, and number of medications taken. Reference value: 11 s.

risk (HR = 0.58, 95 % CI: 0.52, 0.64).

Fig. 5 shows sex-specific analyses of HGS and mortality. In men, the association was steeper, particularly at lower grip strength levels (10th percentile = 32 kg, HR = 1.55, 95 % CI: 1.44, 1.66). In women, the risk increase was slightly less pronounced (10th percentile = 19 kg, HR = 1.42, 95 % CI: 1.31, 1.54).

Taken together, these results demonstrate that both HGS and 5-CST are independently and non-linearly associated with mortality risk, with sex-specific patterns suggesting their complementary roles in clinical screening. A summary of the results is presented in Fig. 6.

Full multivariable Cox regression models, including hazard ratios and 95 % confidence intervals for both 5-CST and HGS, are presented in Online Material.

Discussion

In a large sample of middle-aged and older adults from 28 countries, both 5-CST performance and HGS are gradually associated with the risk of all-cause mortality. Although both tests provide valuable information on mortality risk in older people, HGS is a stronger predictor of all-cause mortality than 5-CST, especially in males. In contrast, the sex-stratified analysis suggests that 5-CST may be especially relevant for identifying women over 50 years of age at high mortality risk, as the increased risk was more sensitive to higher test times in this group. This is the first cohort study with long-term follow-up (mean 7 years) to perform a comparative analysis for the association between both markers and mortality.

Mortality risk of 5-CST

Our results indicate that above 10 s for men and 11 s for women there is an increased risk of mortality. Similar thresholds were recently reported as the optimal cut-off values of the 5-CST for assessing sarcopenia in older adults within the Chinese community.²² Furthermore, our observed percentiles are consistent with EWGSOP2 guidelines,¹⁷ which define poor physical performance as taking ≥15 s to complete the 5-CST. In our cohort, the 50th percentile was 11 s and the 90th percentile was 18 s, placing a substantial proportion of high-risk individuals above this clinical threshold. From previous studies, little is known about mortality risk behaviors above and below this threshold. As a novel finding, our




Fig. 3. Sex-stratified association of the 5-repetition chair-stand test with risk of death during follow-up. Note: The solid line shows the hazard ratios and the dashed lines show the 95 % confidence intervals. Model adjusted for age, sex, body mass index, smoking status, educational level, marital status, geographic region, and number of medications taken.




Fig. 4. Association of Handgrip Strength with risk of death during follow-up in adults older than 50 years ($n = 43,605$). Note: The solid line shows the hazard ratios and the dashed lines show the 95 % confidence intervals. Model adjusted for age, sex, body mass index, smoking status, educational level, marital status, geographic region, and number of medications taken. Reference value: 33 kg.

estimates showed that beyond this point, the slope of mortality risk diverged between sexes as the test time increased, becoming more pronounced in women while stabilizing in men (Fig. 3). This can be explained by few cases among men with slow 5-CST performance, as the confidence interval is quite wide (i.e., increasing the uncertainty).

In addition, sex-related physiological differences may explain this variation, as the mechanisms driving muscle decline and its impact on mortality may differ between men and women. First, the 5-CST, a test designed to assess maximum performance in a short period, is related to functional mobility and muscular power capability (maximum strength generated over a short period of time).^{23,24} Relative muscle power has been reported to be independently and negatively associated with mortality and hospitalization in older adults²⁵ and is more important than strength to prevent falls and predict mortality in older adults.²⁶ Second, men tend to have higher muscle power even in older ages,²⁷ which may help maintain performance and stabilize mortality risk despite slower times. In contrast, women experience more significant age-related muscle atrophy, particularly in type II fibers (faster fibers),²⁸

since from menopause onwards, they experience greater loss of muscle strength than men due to decreased estrogen, which plays a key role in maintaining muscle mass.^{29–31} This hormonal deficiency leads to accelerated muscle atrophy and weakness, which affects the speed of execution in the 5-CST and contributes to a steeper rise in mortality risk after the threshold.

Based on our results and these factors, 5-CST seems more recommendable for measuring mortality risk in women than men. These results highlight the practical utility of the 5-CST as a simple, low-cost, equipment-free tool that can be used in community, primary care and physical therapy settings to identify older women at increased risk of mortality. Its feasibility in a wide range of settings makes it a promising screening tool for targeted interventions in populations with limited access to advanced diagnostic resources. Compared to other functional tests such as gait speed or SPPB, the 5-CST mainly reflects the power of the lower limbs. Given its sensitivity in women, the 5-CST could be incorporated into preventive health assessments to help stratify risk and prioritize early mobility-enhancing interventions.

Mortality risk of HGS

Higher HGS was associated with lower mortality risk in both men and women, though the patterns differed. These findings are consistent with several studies.^{11,32–34} Additionally, a previous study examined all-cause mortality in older adults, finding that HGS up to 42 kg for men and 25 kg for women was associated with lower mortality risk, similar to our study (44 kg for men and 27 kg for women).³⁴ Our 10th percentile values (32 kg for men and 19 kg for women) were slightly above the EWGSOP2 thresholds for low grip strength (<27 kg for men, <16 kg for women),¹⁷ yet already linked to elevated mortality. This reinforces the prognostic relevance of HGS even before reaching clinical cutoffs. Despite previous studies on mortality risk and HGS by sex, little is known about the specific trends of the risk curves above and below the HR = 1. Interestingly, in our results, the mortality risk for men decreases sharply as HGS increases, then tends to stabilize. In contrast, in women, while the relationship is similar, the decline in mortality risk is more gradual, and at higher levels of HGS, the risk remains closer to 1. Also, this can be explained by the few women with high values and confidence interval becoming wider (i.e., increasing the uncertain estimate).

This suggests that although increasing HGS reduces mortality risk in both sexes, the protective effect is more pronounced in men, while in women, the effect is less noticeable. These sex-related differences can be




Fig. 5. Sex-stratified association of handgrip strength with risk of death during follow-up. Note: The solid line shows the hazard ratios and the dashed lines show the 95 % confidence intervals. Model adjusted for age, sex, body mass index, smoking status, educational level, marital status, geographic region, and number of medications taken.




Fig. 6. Summary of main finding including total sample.

explained because, generally, men have greater grip strength than women, and the decline in kg with age differs between the sexes, being faster in men than women,^{35,36} due to the hormonal factors.^{29–31} In conjunction with previous research,¹¹ our findings justify preventive strategies for older adults with low HGS in both sexes. However, although HGS is a valid indicator of overall muscle strength and correlates significantly with lower limb strength, it may not fully reflect specific deficits. Where possible, complementary tests such as isometric knee extension should be considered whenever possible.

Our results, however, consider that the reduction in mortality risk with increased HGS differs between men and women. Therefore, these differences must be considered when assessing HGS and implementing strategies to improve longevity and quality of life.

5-CST and HGS as modifiers of mortality risk

Based on our results, both tools provide valuable information, but HGS is more relevant for identifying those at increased mortality risk in both men and women. Nevertheless, the role of HGS as a modifiable factor after exercise remains controversial.³⁷ A previous meta-analysis showed that resistance training in older adults leads to small but significant improvements in HGS, as reported by various training methods.³⁸ Thus, it is recommended that HGS should not be used to assess overall functional performance after exercise regimens.³⁸ In

contrast, other meta-analyses suggest that exercise interventions increase on average HGS by 3–6 kg compared to usual care, where each 1 kg increase in this test is associated with a 9 % reduction in the risk of all-cause mortality in older adults aged >65 years.^{39,40} The relevance of HGS as a prognostic measure lies in its role as a marker of overall body strength; however, evidence on its association with lower limb strength (e.g., knee extensors) remains conflicting, regardless of age and health status.^{18,19} In contrast, the 5-CST may provide a more specific assessment of strength in the lower limbs and trunk.⁴¹ Low HGS typically indicates general weakness, suggesting a need for overall strength training rather than focusing on isolated muscle groups—except in cases where hand-specific conditions, such as osteoarthritis or rheumatoid arthritis, are present. Another key point is the limited window of opportunity for strength improvement in the HGS compared to the lower limbs, as lower body muscle mass, strength, and power are affected more by aging than are upper body measures.⁴² On the other hand, 5-CST is a simple test that can also be used to train muscle power and, therefore, as a factor to be considered a modifiable risk.^{43–47} Moreover, when using both tests, it is crucial to consider the sex-related risk differences and their sensitivity to change after resistance training programs aimed to reduce mortality risk.^{43,44}

Emerging evidence suggests that these measures may not be interchangeable. For example, Verstraeten et al. demonstrated that in geriatric patients hospitalized for rehabilitation, HGS and CST yielded

substantially different prevalence estimates of probable sarcopenia, with little concordance, and that only HGS - but not CST - was associated with institutionalization and mortality.⁴⁸ Our findings are consistent in showing that HGS is more strongly associated with mortality. However, we also observed that CST had prognostic value, especially in women. This suggests that, while HGS may be preferable for general risk stratification, CST may retain value in specific subgroups or when assessing functional impairment. Another advantage of HGS, beyond its association with mortality, is its reliability and clinical utility in detecting muscle weakness even in cognitively impaired older adults.⁴⁹ Additionally, HGS measured at admission can also predict functional decline in older adults at discharge.⁵⁰

In clinical practice, these findings support the use of HGS and 5-CST not only as predictors of long-term outcomes, but also as entry points for initiating tailored strength and mobility programs. For example, older adults with low HGS may benefit from global resistance training interventions, whereas those with longer 5-CST times may require more targeted exercises to improve lower extremity strength and balance. These stratified approaches could help physical therapists design efficient, individualized programs with a potential impact on survival and quality of life.

Limitations

The results of this study should be interpreted with caution, taking into account the following limitations. First, although the model adjusted for several confounding variables, it may not have been sufficient to completely eliminate confounding factors. In addition, some of these variables may change over time, affecting the relationship between testing and mortality. Thus, there is still the possibility of both residual confounding and confounding over time. Second, as the mortality outcome was obtained through a proxy relative, there is still the possibility of some degree of misclassification bias. Third, since the analytical sample only included participants with valid HGS and 5-CST data, it is possible that people with severe physical or cognitive impairments were excluded, which could introduce a selection bias. Fourth, Although the SHARE protocol standardizes administration across countries, specific details on chair type, safety procedures, or discontinuation rates are not available in the dataset. Similarly, the dataset does not include information on the interval between repetitions of HGS measurements. Finally, although our study sample included individuals in an age range associated with higher mortality, it is plausible that the associations of the chair-stand test extend to younger age groups and require further investigation.

An additional consideration is the geographic diversity of the sample, which included participants from Eastern, Northern, Southern and Western Europe, as well as Israel. This broad representation reinforces the external validity of our results. However, regional differences in health systems and socio-economic conditions may influence associations. Although we adjusted our models for geographical regions, some residual confounding cannot be excluded. Future studies could explore these associations within specific regions to better understand local contextual effects. Furthermore, future research should investigate whether changes in 5-CST and HGS can predict mortality trajectories over time. The effectiveness of interventions aimed at improving these markers in high-risk groups, such as older women, should also be evaluated.

Conclusion

Handgrip strength is a stronger predictor of all-cause mortality than the 5-CST. However, the 5-CST may also be useful, especially in middle-aged and older women. Both tools provide valuable information, but handgrip strength is more relevant for identifying those at increased mortality risk. Given the simplicity and accessibility of both assessments, their implementation as routine screening tools in physical

therapy practice could facilitate early identification of middle-aged and older adults at increased risk of mortality. This would support earlier interventions and better clinical decisions, especially for older adults experiencing functional decline.

Declaration of competing interest

The authors declare no competing interests.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.bjpt.2025.101529.

References

1. Physiological and architectural changes in the ageing muscle and their relation to strength and function in sarcopenia. *Eur Geriatr Med.* 2016;7(3):201–206.
2. Kennis E, Verschueren S, Van Roie E, Thomis M, Lefevre J, Delecluse C. Longitudinal impact of aging on muscle quality in middle-aged men. *Age.* 2014;36(4):9689.
3. Wang DXM, Yao J, Zirek Y, Reijnierse EM, Maier AB. Muscle mass, strength, and physical performance predicting activities of daily living: a meta-analysis. *J Cachexia Sarcopenia Muscle.* 2020;11(1):3–25.
4. United Nations Department of Economic, S Affairs. *World Population Ageing 2019.* United Nations; 2020.
5. Schenkman M, Hughes MA, Samsa G, Studenski S. The relative importance of strength and balance in chair rise by functionally impaired older individuals. *J Am Geriatr Soc.* 1996;44(12):1441–1446.
6. Beaudart C, Zaaria M, Pasleau F, Reginster JY, Bruyère O. Health Outcomes of Sarcopenia: a Systematic Review and Meta-Analysis. *PLoS One.* 2017;12(1), e0169548.
7. Bohannon RW. Muscle strength: clinical and prognostic value of hand-grip dynamometry. *Curr Opin Clin Nutr Metab Care.* 2015;18(5):465–470.
8. Strandkvist V, Larsson A, Pauelsen M, et al. Hand grip strength is strongly associated with lower limb strength but only weakly with postural control in community-dwelling older adults. *Arch Gerontol Geriatr.* 2021;94, 104345.
9. Vaidya SM, Nariya DM. Handgrip strength as a predictor of muscular strength and endurance: a cross-sectional study. *J Clin Diagn Res.* 2021. <https://doi.org/10.7860/jcdr/2021/45573.14437>. Published online.
10. Bohannon RW. Grip strength: an indispensable biomarker for older adults. *Clin Interv Aging.* 2019;14:1681–1691.
11. López-Bueno R, Andersen LL, Koyanagi A, et al. Thresholds of handgrip strength for all-cause, cancer, and cardiovascular mortality: a systematic review with dose-response meta-analysis. *Ageing Res Rev.* 2022;82, 101778.
12. Leong DP, Teo KK, Rangarajan S, et al. Prognostic value of grip strength: findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet.* 2015;386 (9990):266–273.
13. Zhao X, Chen S, Liu N, Hu F, Yu J. Handgrip strength is positively associated with successful aging in older adults: a national cross-sectional study in China. *J Affect Disord.* 2023;333:30–37.
14. Bohannon RW. Reference values for the five-repetition sit-to-stand test: a descriptive meta-analysis of data from elders. *Percept Mot Skills.* 2006;103(1):215–222.
15. Lord SR, Murray SM, Chapman K, Munro B, Tiedemann A. Sit-to-stand performance depends on sensation, speed, balance, and psychological status in addition to strength in older people. *J Gerontol A Biol Sci Med Sci.* 2002;57(8):M539–M543.
16. Buatois S, Perret-Guillaume C, Gueguen R, et al. A simple clinical scale to stratify risk of recurrent falls in community-dwelling adults aged 65 years and older. *Phys Ther.* 2010;90(4):550–560.
17. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing.* 2019;48(1):16–31.
18. Yeung SSY, Reijnierse EM, Trappenburg MC, et al. Handgrip Strength Cannot Be Assumed a Proxy for Overall Muscle Strength. *J Am Med Dir Assoc.* 2018;19(8): 703–709.
19. Tatangelo T, Muollo V, Ghiotto L, Schena F, Rossi AP. Exploring the association between handgrip, lower limb muscle strength, and physical function in older adults: a narrative review. *Exp Gerontol.* 2022;167, 111902.
20. Wave 9. Accessed October 19, 2024. <https://share-eric.eu/data/data-documentation/waves-overview/wave-9>.
21. Börsch-Supan A, Brandt M, Hunkler C, et al. Data Resource Profile: the Survey of Health, Ageing and Retirement in Europe (SHARE). *Int J Epidemiol.* 2013;42(4): 992–1001.
22. Li YH, Wang XH, Ya S, Jiaoling H, Hua N. The optimal cut-off value of five-time chair stand test for assessing sarcopenia among Chinese community-dwelling older adults. *J Cachexia Sarcopenia Muscle.* 2024;15(2):756–764.
23. Sapega AA, Drillings G. The definition and assessment of muscular power. *J Orthop Sports Phys Ther.* 1983;5(1):7–9.
24. Cruz-Montecinos C, Torres-Castro R, Otto-Yáñez M, et al. Which sit-to-stand test best differentiates functional capacity in older people? *Am J Phys Med Rehabil.* 2024;103 (10):925–928.

25. Losa-Reyna J, Alcazar J, Carnicero J, et al. Impact of relative muscle power on hospitalization and all-cause mortality in older adults. *J Gerontol A Biol Sci Med Sci*. 2022;77(4):781–789.
26. Simpkins C, Yang F. Muscle power is more important than strength in preventing falls in community-dwelling older adults. *J Biomech*. 2022;134, 111018.
27. Alcazar J, Aagaard P, Haddock B, et al. Age- and Sex-Specific Changes in Lower-Limb Muscle Power Throughout the Lifespan. *J Gerontol A Biol Sci Med Sci*. 2020;75(7):1369–1378.
28. Nuzzo JL. Sex differences in skeletal muscle fiber types: a meta-analysis. *Clin Anat*. 2024;37(1):81–91.
29. Sipilä S, Taaffe DR, Cheng S, Puolakka J, Toivanen J, Suominen H. Effects of hormone replacement therapy and high-impact physical exercise on skeletal muscle in post-menopausal women: a randomized placebo-controlled study. *Clin Sci (Lond)*. 2001;101(2):147–157.
30. Taaffe DR, Newman AB, Haggerty CL, et al. Estrogen replacement, muscle composition, and physical function: the Health ABC Study. *Med Sci Sports Exerc*. 2005;37(10):1741–1747.
31. Collins BC, Laakkonen EK, Lowe DA. Aging of the musculoskeletal system: how the loss of estrogen impacts muscle strength. *Bone*. 2019;123:137–144.
32. Wang Y, Liu Y, Hu J, et al. Association of handgrip strength with all-cause mortality: a nationally longitudinal cohort study in China. *J Sci Med Sport*. 2022;25(11): 878–883.
33. Jeong W, Moon JY, Kim JH. Association of absolute and relative hand grip strength with all-cause mortality among middle-aged and old-aged people. *BMC Geriatr*. 2023;23(1):321.
34. López-Bueno R, Andersen LL, Calatayud J, et al. Associations of handgrip strength with all-cause and cancer mortality in older adults: a prospective cohort study in 28 countries. *Age Ageing*. 2022;51(5). <https://doi.org/10.1093/ageing/afac117>.
35. Vianna LC, Oliveira RB, Araújo CGS. Age-related decline in handgrip strength differs according to gender. *J Strength Cond Res*. 2007;21(4):1310–1314.
36. Stenholm S, Härkänen T, Sainio P, Heliövaara M, Koskinen S. Long-term changes in handgrip strength in men and women—accounting the effect of right censoring due to death. *J Gerontol A Biol Sci Med Sci*. 2012;67(10):1068–1074.
37. Yaginuma Y, Abe T, Thiebaud RS, Kitamura T, Kawanishi M, Fukunaga T. Can handgrip strength improve following body mass-based lower body exercise? *Biores Open Access*. 2017;6(1):19–27.
38. Labott BK, Bucht H, Morat M, Morat T, Donath L. Effects of exercise training on handgrip strength in older adults: a meta-analytical review. *Gerontology*. 2019;65(6): 686–698.
39. Valenzuela PL, Saco-Ledo G, Morales JS, et al. Effects of physical exercise on physical function in older adults in residential care: a systematic review and network meta-analysis of randomised controlled trials. *Lancet Healthy Longev*. 2023;4(6): e247–e256.
40. Jerez-Roig J, de Brito Macedo Ferreira LM, Torres de Araújo JR, Costa Lima K. Functional decline in nursing home residents: a prognostic study. *PLoS One*. 2017;12(5), e0177353.
41. Silva P, Franco J, Gusmão A, Moura J, Teixeira-Salmela L, Faria C. Trunk strength is associated with sit-to-stand performance in both stroke and healthy subjects. *Eur J Phys Rehabil Med*. 2015;51(6):717–724.
42. Candow DG, Chilibeck PD. Differences in size, strength, and power of upper and lower body muscle groups in young and older men. *J Gerontol A Biol Sci Med Sci*. 2005;60(2):148–156.
43. Kraschewski JL, Sciamanna CN, Poger JM, et al. Is strength training associated with mortality benefits? A 15year cohort study of US older adults. *Prev Med*. 2016; 87:121–127.
44. Saaidifard F, Medina-Inojosa JR, West CP, et al. The association of resistance training with mortality: a systematic review and meta-analysis. *Eur J Prev Cardiol*. 2019;26(15):1647–1665.
45. Lizama-Pérez R, Chiroso-Ríos LJ, Contreras-Díaz G, Jerez-Mayorga D, Jiménez-Lupión D, IJ Chiroso-Ríos. Effect of sit-to-stand-based training on muscle quality in sedentary adults: a randomized controlled trial. *PeerJ*. 2023;11, e15665.
46. Schot PK, Knutzen KM, Poole SM, Mrotek LA. Sit-to-stand performance of older adults following strength training. *Res Q Exerc Sport*. 2003;74(1):1–8.
47. Gonzalez-Bautista E, de Souto, Barreto P, Salinas-Rodriguez A, et al. Clinically meaningful change for the chair stand test: monitoring mobility in integrated care for older people. *J Cachexia Sarcopenia Muscle*. 2022;13(5):2331–2339.
48. Verstraeten LMG, de Haan NJ, Verbeet E, van Wijngaarden JP, Meskers CGM, Maier AB. Handgrip strength rather than chair stand test should be used to diagnose sarcopenia in geriatric rehabilitation inpatients: reSTORing health of acutely unwell adults (RESORT). *Age Ageing*. 2022;51(11). <https://doi.org/10.1093/ageing/afac242>.
49. Garcia PA, Sampaio RX, de Moura JA, de Souza PF, da Costa LB, Dos Santos, Mendes FA. What is the most appropriate handgrip strength testing protocol for sarcopenia screening in older adults with cognitive impairment? *Braz J Phys Ther*. 2024;28(4), 101104.
50. de Moura TG, da Rocha IF, Guedes LS, Alves AT, Garcia PA. Is prehospital physical performance a predictor of functional capacity decline at discharge in hospitalized Brazilian older adults? *Braz J Phys Ther*. 2024;28(1), 100576.