Contents lists available at ScienceDirect



# Brazilian Journal of Physical Therapy



journal homepage: www.elsevier.com/locate/bjpt

# Systematic Review

# The most effective therapeutic exercises for pain intensity in women with fibromyalgia: A systematic review and network meta-analysis<sup> $\star$ </sup>



Álvaro-José Rodríguez-Domínguez <sup>a</sup>, Manuel Rebollo-Salas <sup>b,\*</sup>, Raquel Chillón-Martínez <sup>a</sup>, Abel Rosales-Tristancho <sup>c</sup>, Inmaculada Villa-del-Pino <sup>b</sup>, José-Jesús Jiménez-Rejano <sup>b</sup>

<sup>a</sup> Department of Health and Sports, Pablo de Olavide University, Seville, Spain

<sup>b</sup> Department of Physical Therapy, University of Seville, Seville, Spain

<sup>c</sup> Department of Applied Economics I, University of Seville, Seville, Spain

ARTICLE INFO	A B S T R A C T
ARTICLEINFO Keywords: Fibromyalgia Pain Physical therapy Rehabilitation Therapeutic exercise	Background: Therapeutic exercise (TE) is the only intervention with strong recommendation for fibromyalgia.However, there is controversy as to which type of exercise is the most beneficial.Objective: To determine which TE approach is the most effective in reducing pain intensity in women withfibromyalgia.Methods: A systematic review was performed with a network meta-analysis (NMA). Six databases were searchedfrom inception until January 2024. Randomized controlled trials (RCTs) evaluating the effects of TE on painintensity were included in women with fibromyalgia. Methodological quality was assessed using the Physio-therapy Evidence Database scale. The size of the effect and the clinically important difference were determined inthe short-term (≤3 months) and long-term (>3 months).Results: Sixty-one studies were identified, of which 51 were included in the quantitative synthesis (n = 2873).Fifteen TE interventions and eight comparison interventions (comparators) were identified. Aquatic exercise (p-score: 0.8713) was found to provide best benefits in the short-term and resistance training in the long-term (p-score: 0.9749). Statistically significant differences were found in favor of aquatic exercise, Pilates, qigong,resistance training, dance, functional training, aquatic exercise, virtual reality, and aerobic exercise (in the long-term) compared to usual care.Conclusion: With a moderate level of evidence, our NMA shows that, in the short-term, aquatic exercise is themost effective TE intervention to reduce pain intensity in women with fibromyalgia, while resistance training is the most effective in the long-term. More RCTs are needed to strengthen these findings.

# Introduction

Fibromyalgia is a chronic syndrome characterized by widespread musculoskeletal pain, chronic fatigue, sleep disturbances, and physical disability.<sup>1–4</sup> It is prevalent in 2 to 4 % of the world's population,<sup>2–6</sup> affecting mainly women.<sup>5,7,8</sup> Although it is hypothesized that fibromyalgia is due to a central sensitization process, the etiopathological mechanisms are still unclear.<sup>3,9–12</sup> Studies have identified several muscle abnormalities, such as a reduction in type II fibers and alterations in muscle contraction and metabolism, which can contribute to fatigue and pain in patients with fibromyalgia.<sup>13,14</sup> These findings may explain the

benefits of physical exercise for these patients.

Physical exercise is the only treatment with a strong recommendation for fibromyalgia.<sup>2,3</sup> Several reviews have evaluated the effects of different therapeutic exercise (TE) approaches in patients with fibromyalgia, with benefits observed in almost all of them.<sup>15–22</sup> However, there is controversy as to which approach is the most beneficial. The Cochrane review<sup>23</sup> of 2007 classified aerobic exercise as the "gold standard" for the treatment of fibromyalgia, although in their 2017 review,<sup>18</sup> they concluded that this exercise approach may have little or no long-term effect on pain.

To date, we have not found any publications that compares all TE

 $^{\star}\,$  Registration number: CRD42021293015.

https://doi.org/10.1016/j.bjpt.2025.101226

Received 23 April 2024; Received in revised form 9 August 2024; Accepted 28 April 2025 Available online 3 May 2025

<sup>\*</sup> Corresponding author at: Department of Physical Therapy, University of Seville, C/ Avicena s/n, Seville 41009, Spain. *E-mail address*: mrebollo@us.es (M. Rebollo-Salas).

<sup>1413-3555/© 2025</sup> The Authors. Published by Elsevier España, S.L.U. on behalf of Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

approaches with each other for the treatment of pain in fibromyalgia. Because the disease affects mainly women and the response to treatment is influenced by a multitude of factors, including gender, the objective of this study was to analyze the efficacy of different TE approaches in pain intensity in women with fibromyalgia and to identify which exercise approach is the most effective through a network meta-analysis (NMA) of randomized clinical trials (RCTs).

### Methods

A systematic review with an NMA following the PRISMA-NMA guidelines.<sup>24</sup> The protocol was registered in the PROSPERO database.

Our search was conducted from database inception until January 14, 2024. The databases selected were MEDLINE, CENTRAL, Embase, Web of Science (WoS), Cumulative Index to Nursing and Allied Health Literature (CINAHL), and Scopus. The search strategy and keywords used are given in Supplementary material (Table S.1)

To minimize publication bias, a search was performed on ClinicalTrials.gov. Additional records were searched by hand from relevant literature reviews to supplement the findings of the database.

# Eligibility criteria

The inclusion criteria followed the PICOS (participants, interventions, comparators, outcomes, study design) strategy:

- 1. Type of study: RCTs.
- 2. Participants: adult women diagnosed with fibromyalgia according to the criteria of the American College of Rheumatology for fibromyalgia (ACR 1990/2010/2016).
- Interventions: studies that included any form of TE as the only intervention or a combination of exercises in any of the groups were selected.
- 4. Comparators: any treatment
- 5. Outcomes: pain intensity, evaluated with visual analog scale (VAS 0-10, 0-100)

Trials published in languages other than English, Spanish, French, Italian, or Portuguese were excluded, as were trials that combined ET with other treatments and trials that included men.

#### Data collection process

Two review authors (AR-D, JJ-R) independently performed study selection and data extraction. A third author (MR-S) was consulted in case of disagreement. Studies that did not report the data required for the meta-analysis were excluded from the quantitative synthesis. In general, the data collected were the mean and standard deviation for each treatment and period studied. Given the existence of studies that provided other types of measurements (such as median or quartiles Q1 and Q3), prior estimates were made to obtain an approximation of the mean and standard deviation from the data.<sup>25,26</sup>

A standardized form was used for data extraction, addressing participants, diagnostic criteria, type of intervention, follow-up time, and results obtained (Supplementary material -Table S.2).

#### Risk of bias

The methodological quality was evaluated using the PEDro scale.<sup>27</sup> The quality of the studies was reviewed by two independent evaluators (IV-P, RC-M), and a third evaluator (AR-T) was consulted when discrepancies appeared. The included studies were classified according to scores of 9 or 10, 6 to 8, and  $\leq$  5 on the PEDro scale and were interpreted as excellent, good, and fair quality, respectively.<sup>28</sup>

# Certainty of the evidence

The CINeMA web application was used to assess confidence in findings from primary NMA. The CINeMA framework considers six domains that affect the level of confidence in the NMA results: (a) withinstudy bias, (b) reporting bias, (c) indirectness, (d) imprecision, (e) heterogeneity, and (f) incoherence. The reviewers assessed the level of concern for each relative treatment effect of NMA as giving rise to "no concerns", "some concerns", or "major concerns" in each of the six domains. Then, judgments across the domains are summarized into a single confidence rating ("high", "moderate", "low", or "very low").<sup>29</sup> (Supplementary material -Table S.3)

# Clinically important differences (CID)

According to Cochrane,<sup>16</sup> the Initiative on Methods, Measurement and Pain Assessment in Clinical Trials (IMMPACT) recommended the following benchmarks for interpreting changes in pain intensity on a numerical rating scale 0 to 10 in chronic pain clinical trials: a) a 10 to 20 % decrease is minimally important, b) a decrease greater than 30 % is moderately important, and c) a decrease greater than 50 % is substantial.<sup>30,31</sup> Therefore, the minimum CID in outcome was interpreted as a pain difference of 15 points out of 100 (VAS 0–100)<del>.</del> The CID was determined from the difference obtained for each intervention compared to usual care.

# Summary measures

The analysis included a qualitative synthesis (Supplementary material -Table S.2) and a quantitative synthesis (pairwise meta-analysis and NMA). VAS is reported as a continuous measure from 0 to 10 and from 0 to 100. The outcome data was standardized from 0–100, using the mean difference (MD) to perform the primary analyzes. A temporal division was established into two periods: short-term ( $\leq$  3 months) and long-term (> 3 months). For the short term, only follow-up results closer to 3 months after the end of the intervention were selected. For the long term, only follow-up results closer to six months after the end of the intervention were selected.

In those meta-analyses, a random-effects model was used where heterogeneity between studies was observed, while a fixed-effects model was used for those where homogeneity was observed. As indicative of homogeneity, I<sup>2</sup> coefficient values <50 % and/or Chi<sup>2</sup> test values of homogeneity with p > 0.05 were taken. In all cases, the corresponding forest plot is presented (Supplementary material -Figs. 2–14). A pairwise meta-analysis was performed using Review Manager software version 5.4.

Regarding NMA, the means provided by each study were taken for each of the treatments performed, as well as the standard deviation of each one, which allowed the calculation of the MD, as well as its standard error.<sup>32</sup> The NMA allows us to know the effect size between any pair of treatments, whether they have been compared directly or not. To establish a ranking between all treatments, it is possible to assign a p-score to each treatment. This score is based on the probability that a treatment is better than any other treatment included in the NMA.<sup>33–35</sup>

# Planned methods of analysis

The NMA developed in this study is framed within the framework of frequentist statistics.  $^{36,37}$  NMA is performed in RStudio using the 'netmeta' library.  $^{38}$ 

Coherence is the statistical tool to test the transitivity between treatment effects.<sup>33</sup> The objective is to study whether there are statistically significant differences between direct and indirect comparisons of any two treatments. For this purpose, the SIDE (Separating Indirect from Direct Evidence) method is used to study local coherence and the I<sup>2</sup> statistic is used in the case of global coherence.<sup>33,39,40</sup> This statistic is

# calculated from Cochran's Q.40

The NMA is developed according to a random-effects model that considers the heterogeneity of the estimates. Furthermore, the random-effects model facilitates extrapolation of the results obtained to a larger population.  $^{39,41}$ 

# Results

Fig. 1 describes the study selection process. Sixty articles were included in the qualitative synthesis<sup>42-101</sup> and 51 were included in the quantitative synthesis (Supplementary material -Table S.2).

# Characteristics of the study and results of individual studies

All studies were published between 1992 and 2021. Three studies<sup>102–104</sup> had two treatment arms with the same exercise approach but different doses, so they were counted as six studies in quantitative synthesis.

The sample size in each article was between 15 and 207, with an average of 59 participants. The total number of participants was 3581, of which 2873 were included in the quantitative synthesis (708 not included). The mean age was approximately  $48.9 \pm 6.1$  years (range = 32–59 years). Of the 60 studies included, 48 used the ACR1990 criteria, 13 the ACR2010 criteria, and 2 the ACR2016 criteria. The total number of studies and participants per exercise modality, the total number of



Fig. 1. Flow chart of the selection of studies.

participants per nationality and intervention method, and the diagnostic criteria used are detailed in Table S.4 (Supplementary material).

# Intervention group (Therapeutic exercise)

The interventions were classified according to the definitions given in Table S.5 (Supplementary material) and described according to the exercise approach used, the program design, and the dosage (intensity, frequency, and duration) (Supplementary material -Table S.2). - One of the identified categories (vibration exercise) was only investigated in one study.<sup>105</sup> However, it did not provide the necessary data to be included in the meta-analysis, so it was only included in the qualitative synthesis<del>.</del> Thus, of the 15 TE categories identified, 14 were included in the NMA.

#### Comparison group (comparator intervention)

Eight intervention categories were identified as comparison groups in the included studies (Supplementary material -Table S.5). Studies that combined exercise therapy with another treatment approach were



**Fig. 2.** Network plot presenting the trial data contributing evidence comparing exercise treatment types. A: Short-term ( $\leq$ 3 months): 37 trials, 171 comparisons, 1922 participants. B: Long-term (>3 months): 24 trials, 136 comparisons, 1530 participants. The size of the nodes represents how many times the exercise appears in any comparison about that treatment and the width of the edges represents the total sample size in the comparisons it connects.

classified as "Combined Therapy" and therefore qualified as a comparison group. Studies that included a placebo treatment were classified as "sham" and those in which no intervention or usual care was performed as "usual care".

#### Outcomes

The main analyzes of this study evaluated the effect of the different identified TE approaches on pain intensity in women with fibromyalgia, assessed with VAS. All available results were collected for the available follow-up time points, with two-time cut-off points for the primary meta-analyzes: short-term ( $\leq$ 3 months) and long-term (>3 months).

# Synthesis of results

The network formed by the treatments and their comparisons is

constructed from a graph where each node represents a treatment and each edge a direct comparison.<sup>106</sup> The size of the nodes is proportional to the sample collected for each treatment. However, the thickness of each edge is proportional to the number of comparisons collected between the two treatments at its ends.

# Pain results

For the short-term (Fig. 2A), a total of 19 interventions were analyzed by 37 different studies (35 studies plus two studies that had two treatment arms of the same intervention). Among them, there were 32 pairwise comparisons and five comparisons between the three intervention groups at the same time. Usual care had the largest sample size (n = 307), followed by mixed exercise (n = 290), aquatic exercise (n = 236), flexibility (n = 234), and aerobic exercise (n = 233). The thickness of the edges determines that there are a greater number of





Fig. 3. Treatment rankings and surface under the cumulative ranking curve (SUCRA) for pain intensity. A: Short-term ( $\leq$ 3 months); B: Long-term (>3 months).

comparisons between aquatic exercise and mixed exercise (4 comparisons) and between flexibility and resistance training (3 comparisons).

Regarding transitivity, the SIDE method did not show statistically significant differences between direct and indirect comparisons of treatments (Supplementary material -Table S.6A). Furthermore, the  $I^2$  statistic had a value of 84.7 % (78.5 %, 89.1 %).

Fig. 3 shows the ranking between all treatments. The p-score represented for each TE intervention in Fig. 3A places the aquatic exercise group in the first position (with a value of 0.8713), followed by Pilates (0.775).

Fig. 4 shows the relative effects between all pairs of comparisons and their respective confidence intervals at the 95 % level. A negative value of the effect size indicates the superiority of the column treatment, while a positive value indicates the superiority of the row treatment. The order of the columns in Fig. 4A was constructed following the ranking established in Fig. 3A.

Statistically significant differences were found in favor of aquatic exercise, Pilates, qigong, resistance training, virtual reality, mixed exercise, and aerobic exercise compared to usual care and sham. Aquatic exercise also obtained statistically significant differences in comparisons with mixed exercise, aerobic exercise, flexibility, and core stability. Taking the usual care as a reference, it is possible to observe the effect size with respect to the other interventions in Fig. 5A.

Finally, 12 short-term pairwise meta-analyses were also performed (Supplementary material -Figs. S1-S6 and S.9-S14), of which five TE approaches showed statistically significant results in favor of aquatic exercise (p < 0.0001), Tai Chi (p < 0.00001), virtual reality (p = 0.001), resistance training (p = 0.02), and qigong (p = 0.02). On the other hand, an intervention (flexibility) had statistically significant differences in favor of the comparison group, with a p-value <0.005 (p = 0.002). All results are summarized in Table S. 7 (Supplementary material).

For the long-term (Fig. 2B), there were 17 interventions evaluated by 24 studies (22 studies, and two more with one pair of treatment arms each). Of the 24 studies, 20 made pairwise comparisons and four presented comparisons between three treatments at the same time. Node thickness again indicates that the usual care group has the largest sample size (n = 312), followed by the aerobic exercise (n = 177), mixed exercise (n = 164), and aquatic exercise (n = 156) groups. Furthermore, the number of comparisons between the usual care and aerobic exercise groups was the highest, with a total of four between all studies, which resulted in a higher thickness on the corresponding edge in Fig. 2B.

Regarding transitivity, the net splitting method also did not show statistically significant differences between direct and indirect comparisons of treatments (Supplementary material -Table S.6B). Furthermore, the  $I^2$  statistic had a value of 0 % (0 %, 56.6 %).

The p-score represented for each TE intervention in Fig. 3B places resistance training first (0.9749), followed by dance (0.905) and functional training (0.8502).

The order of the columns in Fig. 4B was also constructed following the ranking established in Fig. 3B.Fig. 4B shows that a greater number of statistically significant comparisons were obtained in the long term than in the short term. Thus, we can highlight that statistically significant differences were found in favor of resistance training, dance, functional training, aquatic exercise, virtual reality, and aerobic exercise compared to usual care, and statistically significant differences were found in favor of resistance training, aquatic exercise, aerobic exercise, and mixed exercise compared to education. Furthermore, resistance training showed statistically significant differences in all comparisons, except compared to dance and functional training. Fig. 5B shows the effect size using usual care as a reference group.

Finally, 10 pairwise meta-analyses were performed in the long-term (Supplementary material -Figs. S.1-S.9 and S.13), of which six TE approaches showed statistically significant results in their favor: resistance training (p < 0.0001), aquatic exercise (p = 0.0002), dance (p = 0.003), functional training (p = 0.004), aerobic exercise (p = 0.008), and virtual reality (p = 0.04). As in the short-term, flexibility had statistically

significant differences in favor of the comparison group, with a p-value  $<0.005 \ (p = 0.003)$ . All results are summarized in Table S.7 (Supplementary material).

#### Clinically important differences (CID)

Fig. 5 shows the CID of the interventions analyzed compared to usual care. In the short-term (Fig. 4A), CIDs were obtained for many TE approaches (aquatic exercise, Pilates, qigong, resistance training, virtual reality, mixed exercise, Tai Chi, and aerobic exercise). However, in the long term (Fig. 4B), only three interventions achieved a CID: resistance training, dance, and functional training. Furthermore, only resistance training showed a moderately important clinical difference ( $\geq$ 30 points out of 100). Finally, in Fig. 4A and 4B, some comparisons between TE approaches and the comparators presented CID in favor of the comparators.

# Risk of bias

Most studies (48 out of 60) were rated as "good quality" on the PEDro scale. Seven studies were rated as "excellent quality" and five studies were rated as "fair quality" (Supplementary material -Table S.8).

# Publication bias risk assessment

The Begg and Egger tests did not reveal statistical evidence of publication bias (p > 0.05). These findings are shown in the funnel plots (Supplementary material -Fig. S.15-S.28). The sensitivity analysis indicated that the general results were not substantially modified by eliminating any result.

### Certainty of the evidence

In the short-term, 171 comparisons (47 studies) were performed, of which 22 were rated as "moderate quality". The remaining 149 comparisons were rated as "low quality". Long-term, 137 comparisons (32 studies) were performed, of which only one was rated as "high quality" (aerobic exercise versus usual care). Of the remaining 136 comparisons, 110 were rated as "moderate quality" and 26 as "low quality". (Supplementary material -Table S.98).

# Discussion

To our knowledge, this is the first NMA to compare different TE approaches used in women with fibromyalgia to reduce pain intensity. Our NMA established a ranking (Fig. 3) of TE approaches that we used to order the columns in Fig. 4 showing the relative effects between all pairs of comparisons. Therefore, Fig. 4 can be used as a valuable decision-making tool when choosing the most appropriate type of TE to reduce pain intensity in women with fibromyalgia.

In the short-term (Fig. 4A), we had seven TE approaches that generated statistically significant differences compared to usual care and sham; also, these improvements were clinically important. If we analyze the comparisons between these seven approaches, our results appear to indicate that, in the short term, the five most effective type of TE to improve pain intensity in women with fibromyalgia were aquatic exercise, Pilates, qigong, resistance training, and virtual reality. Furthermore, the strength of evidence, in all these comparisons with usual care was moderate, except for Pilates, which was low.

In the long-term (Fig. 4B), we had six TE approaches that generated statistically significant differences compared to usual care and education. Compared to sham, five types of TE generated statistically significant differences. If we analyze the comparisons between these exercise approaches, our results appear to indicate that, in the long-term, the three most effective type of TE, compared to usual care, education, and sham, to improve pain intensity in women with fibromyalgia, were -

												Aquatic Exercise
											Pilates	-2.80 (-21.16, 15.57)
	Α									Qigong	-4.86 (-31.13, 21.41)	-7.65 (-27.75, 12.44)
									Resistance Training	-2.17 (-25.36, 21.01)	-7.03 (-29.26, 15.19)	-9.83 (-25.76, 6.11)
								Virtual Reality	-1.27 (-21.08, 18.54)	-3.45 (-26.67, 19.78)	-8.30 (-31.58, 14.98)	-11.10 (-27.77, 5.57)
							Mixed Exercise	-0.38 (-15.84, 15.07)	-1.65 (-17.50, 14.19)	-3.83 (-22.58, 14.92)	-8.69 (-28.32, 10.94)	-11.48 (-21.06, -1.90)
						Tai Chi	-0.59 (-24.88, 23.71)	-0.97 (-28.04, 26.10)	-2.24 (-29.65, 25.16)	-4.42 (-33.56, 24.73)	-9.27 (-39.43, 20.88)	-12.07 (-37.10, 12.96)
					Aerobic Exercise	-0.05 (-24.81, 24.72)	-0.63 (-10.94, 9.68)	-1.02 (-17.75, 15.71)	-2.29 (-17.34, 12.76)	-4.46 (-23.84, 14.91)	-9.32 (-29.30, 10.66)	-12.12 (-23.03, -1.20)
				Yoga	-6.75 (-33.98, 20.48)	-6.80 (-40.17, 26.57)	-7.39 (-34.19, 19.42)	-7.77 (-37.12, 21.58)	-9.04 (-38.69, 20.61)	-11.22 (-42.49, 20.06)	-16.07 (-48.28, 16.14)	-18.87 (-46.34, 8.61)
			Flexibility	0.14 (-27.52, 27.80)	-6.61 (-17.78, 4.55)	-6.66 (-31.90, 18.58)	-7.24 (-18.36, 3.87)	-7.63 (-23.49, 8.24)	-8.90 (-22.35, 4.55)	-11.07 (-31.60, 9.45)	-15.93 (-34.23, 2.37)	-18.73 (-29.56, -7.90)
		Core Stability	-4.00 (-26.50, 18.51)	-3.86 (-35.82, 28.11)	-10.61 (-32.63, 11.41)	-10.66 (-40.55, 19.24)	-11.24 (-32.09, 9.61)	-11.63 (-36.30, 13.05)	-12.90 (-37.90, 12.10)	-15.07 (-41.99, 11.84)	-19.93 (-47.77, 7.92)	-22.72 (-44.84, -0.61)
	Balance Exercise	-1.72 (-36.52, 33.08)	-5.72 (-34.89, 23.44)	-5.58 (-43.89, 32.73)	-12.33 (-39.28, 14.61)	-12.38 (-48.98, 24.21)	-12.97 (-41.82, 15.88)	-13.35 (-45.07, 18.37)	-14.62 (-45.49, 16.24)	-16.80 (-49.98, 16.39)	-21.65 (-55.20, 11.89)	-24.45 (-53.52, 4.62)
Usual Care	-4.62 (-33.72, 24.48)	-6.34 (-26.37, 13.68)	-10.34 (-22.36, 1.68)	-10.20 (-35.12, 14.72)	-16.95 (-27.94, -5.97)	-17.00 (-39.19, 5.19)	-17.59 (-27.47, -7.70)	-17.97 (-33.48, -2.46)	-19.24 (-35.32, -3.16)	-21.42 (-40.31, -2.52)	-26.27 (-46.69, -5.86)	-29.07 (-40.65, -17.49)
Sham	-13.85 (-47.29, 19.60)	-15.57 (-44.21, 13.06)	-19.57 (-41.56, 2.42)	-19.43 (-52.22, 13.36)	-26.18 (-45.99, -6.37)	-26.23 (-57.00, 4.54)	-26.82 (-47.88, -5.75)	-27.20 (-52.16, -2.24)	-28.47 (-52.80, -4.15)	-30.64 (-51.84, -9.45)	-35.50 (-63.00, -8.01)	-38.30 (-60.05,- 16.54)
Acupuncture	2.60 (-34.15, 39.36)	0.88 (-21.98, 23.74)	-3.12 (-28.57, 22.33)	-2.98 (-36.79, 30.84)	-9.73 (-34.73, 15.27)	-9.78 (-41.64, 22.08)	-10.36 (-34.62, 13.89)	-10.75 (-38.07, 16.58)	-12.02 (-39.66, 15.62)	-14.19 (-43.57, 15.18)	-19.05 (-49.35, 11.25)	-21.85 (-47.02, 3.33)
Relaxation	4.22 (-35.24, 43.68)	2.50 (-32.57, 37.56)	-1.50 (-29.52, 26.52)	-1.36 (-39.88, 37.16)	-8.11 (-36.94, 20.71)	-8.16 (-44.98, 28.66)	-8.75 (-37.99, 20.50)	-9.13 (-40.70, 22.44)	-10.40 (-34.98, 14.18)	-12.57 (-46.37, 21.22)	-17.43 (-50.57, 15.71)	-20.23 (-49.52, 9.07)
Drugs Therapy	15.19 (-19.83, 50.21)	13.46 (-17.50, 44.43)	9.47 (-15.23, 34.16)	9.61 (-25.25, 44.46)	2.85 (-19.51, 25.22)	2.81 (-30.16, 35.77)	2.22 (-21.89, 26.33)	1.84 (-25.69, 29.36)	0.57 (-26.16, 27.29)	-1.61 (-28.62, 25.41)	-6.47 (-36.18, 23.25)	-9.26 (-33.79, 15.27)
Manual Therapy	17.85 (-23.58, 59.29)	16.13 (-20.85, 53.10)	12.13 (-18.31, 42.57)	12.27 (-28.09, 52.63)	5.52 (-25.96, 37.00)	5.47 (-33.27, 44.21)	4.89 (-26.37, 36.14)	4.50 (-29.17, 38.17)	3.23 (-29.72, 36.18)	1.06 (-34.74, 36.86)	-3.80 (-28.12, 20.52)	-6.60 (-37.07, 23.88)
Combined Therapy	22.99 (-10.24, 56.22)	21.27 ( -0.24, 42.78)	17.27 ( -2.65, 37.19)	17.41 (-13.70, 48.52)	10.66 ( -8.79, 30.11)	10.61 (-18.37, 39.59)	10.03 ( -6.97, 27.02)	9.64 (-12.91, 32.19)	8.37 (-14.47, 31.21)	6.20 (-18.74, 31.14)	1.34 (-24.37, 27.05)	-1.46 (-20.68, 17.77)

Resistance Training											
-9.88 (-31.16, 11.40)	Dance										
-13.20 (-38.72, 12.32)	-3.32 (-20.40, 13.76)	Functional Training								R	
-24.97 (-45.85, -4.09)	-15.09 (-23.84, -6.34)	-11.77 (-28.35, 4.81)	Aquatic Exercise							D	
-26.68 (-48.71, -4.65)	-16.80 (-28.02, -5.59)	-13.48 (-31.49, 4.53)	-1.71 (-12.15, 8.7	73) Virtual Reality							
-27.53 (-54.80, -0.27)	-17.66 (-37.25, 1.93)	-14.33 (-38.46, 9.80)	-2.56 (-21.19, 16.07)	-0.85 (-21.26, 19.55)	Pila	tes					
-28.73 (-49.62, -7.84)	-18.86 (-27.63, -10.08)	-15.53 (-32.13, 1.06)	-3.76 (-10.11, 2.5	58) -2.05 (-12.52, 8.41)	-1.20 (-1	18.72, 2)	Aerobic Exercise				
-30.60 (-52.78, -8.42)	-20.72 (-32.24, -9.21)	-17.40 (-35.60, 0.80)	-5.63 (-16.39, 5.1	-3.92 (-16.77, 8.93)	-3.07 (-2 17.5	23.64, 0)	-1.87 (-12.65, 8.92)	Core Stability			
-33.33 (-56.01, -10.65)	-23.45 (-35.89, -11.01)	-20.13 (-38.92, -1.34)	-8.36 (-17.20, 0.4	48) -6.65 (-20.33, 7.03)	-5.80 (-2 14.8	26.42, 3)	-4.60 (-15.48, 6.29)	-2.73 (-16.66, 11.20)	Mixed Exercise		
-40.53 (-64.55, -16.52)	-30.66 (-45.39, -15.92)	-27.33 (-47.72, -6.95)	-15.56 (-29.00, -2.13)	-13.85 (-29.66, 1.95)	, -13.00 (-25 -0.09)		-11.80 (-23.64, 0.04)	-9.93 (-25.95, 6.08)	-7.20 (-23.29, 8.88)	Flexibility	
-36.00 (-56.15, -15.85)	-26.12 (-32.95, -19.29)	-22.80 (-38.46, -7.14)	-11.03 (-16.50, -5.56)	-9.32 (-18.22, -0.42)	-8.47 (-26.83, 9.90)		-7.27 (-12.77, -1.76)	-5.40 (-14.67, 3.87)	-2.67 (-13.07, 7.73)	4.53 ( -8.52, 17.59)	Usual Care
-41.36 (-64.82, -17.89)	-31.48 (-45.30, -17.66)	-28.16 (-47.89, -8.42)	-16.39 (-27.08, -5.69)	-14.68 (-29.63, 0.27)	-13.82 (-35.31, 7.66)		-12.62 (-25.06, -0.19)	-10.76 (-25.93, 4.42)	-8.03 (-14.05, -2.00)	-0.82 (-17.99, 16.35)	Education
-39.63 (-63.09, -16.18)	-29.76 (-43.56, -15.95)	-26.43 (-46.16, -6.71)	-14.66 (-27.07, -2.26)	-12.95 (-27.89, 1.98)	-12.10 (-32.61, 8.41)		-10.90 (-21.56, -0.24)	-9.03 (-24.20, 6.13)	-6.30 (-21.54, 8.93)	0.90 (-15.03, 16.83)	Sham
-31.33 (-54.46, -8.21)	-21.46 (-34.69, -8.22)	-18.13 (-37.46, 1.19)	-6.36 (-18.13, 5.4	40) -4.65 (-19.06, 9.76)	-3.80 (-23.93, 16.33)		-2.60 (-12.51, 7.31)	-0.73 (-15.38, 13.91)	2.00 (-12.72, 16.71)	9.20 ( -6.24, 24.64)	Drugs Therapy
-30.40 (-52.42, -8.38)	-20.52 (-31.71, -9.33)	-17.20 (-35.19, 0.79)	-5.43 (-15.84, 4.9	-3.72 (-16.28, 8.84)	-2.87 (-23.26, 17.52)		-1.67 (-12.10, 8.77)	0.20 ( -9.09, 9.49)	2.93 (-10.73, 16.59)	10.13 ( -5.65, 25.92)	Acupunctre
-30.62 (-54.05, -7.20)	-20.75 (-34.50, -6.99)	-17.42 (-37.11, 2.27)	-5.65 (-16.27, 4.9	-3.94 (-18.83, 10.95)	-3.09 (-24.53, 18.35)		-1.89 (-14.25, 10.48)	-0.02 (-15.14, 15.09)	2.71 ( -3.16, 8.58)	9.91 ( -7.21, 27.03)	Combined Therapy
-15.65 (-22.21, -9.08)	-5.77 (-28.04, 16.50)	-2.45 (-28.80, 23.90)	9.32 (-12.57, 31.2	21) 11.03 (-11.96, 34.02)	11.88 (-16.16, 39.93)		13.08 ( -8.81, 34.98)	14.95 ( -8.18, 38.09)	17.68 ( -5.93, 41.29)	24.88 ( -0.01, 49.78)	Relaxation
	Statistically significant and clinically important difference favouring column treatment			Clinically important dife favouring column treat	errence tment	Statistica favour	ally significant different ring column treatment	nce No difference	Clinically impor favouring ro	rtant difference w treatment	

Fig. 4. League table of network meta-analysis results for all comparisons between exercise and non-exercise interventions. Effects are expressed as the mean difference (95 % CI) between interventions. A: short-term outcomes ( $\leq$ 3 months); B: long-term results (>3 months).



Fig. 5. Summary network meta-analysis results for each exercise type compared with Usual Care. A: short-term ( $\leq$ 3 months); B: long term (>3 months). The hashed line indicates clinically important difference.

resistance training, dance, and functional training. Furthermore, resistance training and dance were the most effective TE compared to drug therapy, acupuncture, and combination therapy; and only resistance training was the most effective TE compared to relaxation. All of these improvements were clinically important. The strength of evidence, of all these comparisons was moderate.

Importantly, resistance training was the only intervention that showed statistically significant and clinically relevant differences compared to usual care and sham in both the short and long term. It was also the only intervention that showed a moderately important clinical difference ( $\geq$ 30 points out of 100), so, overall, resistance training may be the most effective type of TE to reduce pain intensity in women with fibromyalgia.

Our findings have significant clinical implications for several reasons. Most importantly, Fig. 4 can serve as a valuable guide for selecting the most appropriate TE approaches to reduce pain intensity in women with fibromyalgia. This decision should also be made based on the patient's characteristics and preferences, as this is a key factor in improving adherence to treatment.<sup>107–109</sup> These findings could also contribute to a change in the current paradigm regarding TE prescription in fibromyalgia, because the "gold standard" recommendation of aerobic exercise should be replaced by other more effective approaches. Specifically, aquatic exercise was more effective than aerobic exercise in the short-term and resistance training and dance were more effective in the long-term.

This review has several limitations. First, studies are at risk of bias due to the impossibility of blinding when active interventions are used. Second, this review is limited to the effects of exercise in women with fibromyalgia. It seems more appropriate to limit the sample to this sex, as fibromyalgia affects mainly women and including men in the review would increase the risk of bias. Therefore, these results cannot be extrapolated to men with fibromyalgia. Third, there are concerns about the patient selection criteria used in RCTs. A total of 80 % of the trials used the ACR 1990 criteria, the validity of which was refuted by the authors themselves in 2010. Despite this, studies published in 2021 still continue to use these criteria.<sup>110–112</sup> This could lead to serious selection bias in patients, which could affect the extrapolation of the results to the clinical setting. Finally, our study has evaluated the efficacy of different TE approaches in reducing pain intensity. However, it should be noted that intensity is only one dimension of pain experience, and therefore an

improvement in pain intensity does not necessarily imply an improvement in the overall impact of the disease.  $^{19}\,$ 

Future studies should focus on whether different parameters of the FITT model (frequency, intensity, time, and type), could optimize outcomes in reducing pain intensity in women with fibromyalgia. Finally, it is imperative to unify diagnostic criteria in fibromyalgia so that these patients are classified more accurately and homogeneously. The criteria currently used for the diagnosis of fibromyalgia are still based on a biomedical model, ignoring the recommendations of the World Health Organisation (WHO). Psychosocial variables are known to have a significant impact on disease development, so it is necessary to establish diagnostic criteria based on a biopsychosocial model.<sup>113</sup>

# Conclusions

The NMA showed, with a moderate level of evidence, that the most effective TE approach to reduce pain intensity in women with fibromyalgia was, in the short-term, aquatic exercise and, in the long-term, resistance training. Furthermore, resistance training was the only intervention that showed short- and long-term improvements, with a moderately important clinical difference. More RCTs are needed to strengthen these findings.

# Declaration of competing interest

The authors declare no competing interest.

#### Acknowledgements

We appreciate the work of all the authors who conducted the clinical trials that were included in our meta-analysis.

# Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.bjpt.2025.101226.

# References

<sup>1.</sup> Andrade A, Dominski FH, Sieczkowska SM. What we already know about the effects of exercise in patients with fibromyalgia: an umbrella review. *Semin* 

Arthritis Rheum. 2020;50(6):1465–1480. https://doi.org/10.1016/J. SEMARTHRIT.2020.02.003.

- Macfarlane GJ, Kronisch C, Dean LE, et al. EULAR revised recommendations for the management of fibromyalgia. *Ann Rheum Dis.* 2017;76(2):318–328. https:// doi.org/10.1136/ANNRHEUMDIS-2016-209724.
- Sarzi-Puttini P, Giorgi V, Marotto D, Atzeni F. Fibromyalgia: an update on clinical characteristics, aetiopathogenesis and treatment. *Nat Rev Rheumatol.* 2020;16(11): 645–660. https://doi.org/10.1038/s41584-020-00506-w, 2020 16:11.
- Marques AP, Santo A de S do E, Berssaneti AA, Matsutani LA, Yuan SLK. Prevalence of fibromyalgia: literature review update. *Rev Bras Reumatol.* 2017;57(4):356–363. https://doi.org/10.1016/J.RBRE.2017.01.005.
- Cabo Meseguer A, Cerdá-Olmedo G, Trillo Mata JL. Fibromyalgia: prevalence, epidemiologic profiles and economic costs. *Med Clin.* 2017;149(10):441–448.
- Galvez-Sánchez CM, Duschek S, Del Paso GAR. Psychological impact of fibromyalgia: current perspectives. Psychol Res Behav Manag. 2019;12:117. https://doi.org/10.2147/PRBM.S178240.
- Wolfe F, Walitt B, Perrot S, Rasker JJ, Häuser W. Fibromyalgia diagnosis and biased assessment: sex, prevalence and bias. *PLoS One*. 2018;13(9), e0203755.
- Walitt B, Katz RS, Bergman MJ, Wolfe F. Three-quarters of persons in the US population reporting a clinical diagnosis of fibromyalgia do not satisfy fibromyalgia criteria: the 2012 National Health Interview Survey. *PLoS One*. 2016; 11(6), e0157235.
- Silverwood V, Chew-Graham CA, Raybould I, Thomas B, Peters S. If it's a medical issue i would have covered it by now": learning about fibromyalgia through the hidden curriculum: a qualitative study. *BMC Med Educ.* 2017;17(1):1–9. https:// doi.org/10.1186/S12909-017-0972-6/TABLES/1.
- Yunus MB. Central Sensitivity syndromes: a new paradigm and group nosology for fibromyalgia and overlapping conditions, and the related issue of disease versus illness. *Semin Arthritis Rheum.* 2008;37(6):339–352. https://doi.org/10.1016/J. SEMARTHRIT.2007.09.003.
- Cagnie B, Coppieters I, Denecker S, Six J, Danneels L, Meeus M. Central sensitization in fibromyalgia? A systematic review on structural and functional brain MRI. Semin Arthritis Rheum. 2014;44(1):68–75. https://doi.org/10.1016/J. SEMARTHRIT.2014.01.001.
- Meeus M, Nijs J. Central sensitization: a biopsychosocial explanation for chronic widespread pain in patients with fibromyalgia and chronic fatigue syndrome. *Clin Rheumatol.* 2006;26(4):465–473.
- Bidonde J, Busch AJ, Webber SC, et al. Aquatic exercise training for fibromyalgia. Cochrane Database Syst Rev. 2014;(10). https://doi.org/10.1002/14651858. CD011336/MEDIA/CDSR/CD011336/IMAGE\_N/NCD011336-CMP-012-02.PNG, 2014.
- Park JH, Niermann KJ, Olsen N. Evidence for metabolic abnormalities in the muscles of patients with fibromyalgia. *Curr Rheumatol Rep 2000 2:2.* 2000;2(2): 131–140.
- Häuser W, Klose P, Langhorst J, et al. Efficacy of different types of aerobic exercise in fibromyalgia syndrome: a systematic review and meta-analysis of randomised controlled trials. *Arthritis Res Ther.* 2010;12(3):1–14. https://doi.org/10.1186/ AR3002/TABLES/5.
- Busch AJ, Webber SC, Richards RS, et al. Resistance exercise training for fibromyalgia. *Cochrane Database Syst Rev.* 2013;(12). https://doi.org/10.1002/ 14651858.CD010884/MEDIA/CDSR/CD010884/IMAGE\_N/NCD010884-CMP-006-06.PNG, 2013.
- Nelson NL. Muscle strengthening activities and fibromyalgia: a review of pain and strength outcomes. J Bodyw Mov Ther. 2015;19(2):370–376. https://doi.org/ 10.1016/J.JBMT.2014.08.007.
- Bidonde J, Busch AJ, Schachter CL, et al. Aerobic exercise training for adults with fibromyalgia. *Cochrane Database System Rev.* 2017;2017(6). https://doi.org/ 10.1002/14651858.CD012700/MEDIA/CDSR/CD012700/IMAGE\_N/NCD012700-CMP-004-06.PNG.
- Sosa-Reina MD, Nunez-Nagy S, Gallego-Izquierdo T, Pecos-Martín D, Monserrat J, Álvarez-Mon M. Effectiveness of therapeutic exercise in Fibromyalgia syndrome: a systematic review and meta-analysis of randomized clinical trials. *Biomed Res Int.* 2017. https://doi.org/10.1155/2017/2356346, 2017.
- Andrade A, de Azevedo Klumb Steffens R, Sieczkowska SM, Peyré Tartaruga LA, Torres Vilarino G. A systematic review of the effects of strength training in patients with fibromyalgia: clinical outcomes and design considerations. *Adv Rheumatol.* 2019;58(1):36. https://doi.org/10.1186/S42358-018-0033-9.
- Bidonde J, Busch AJ, Schachter CL, et al. Mixed exercise training for adults with fibromyalgia. *Cochrane Database Syst Rev.* 2019;2019(5). https://doi.org/10.1002/ 14651858.CD013340/MEDIA/CDSR/CD013340/IMAGE\_N/NCD013340-CMP-004-06.PNG.
- Bravo C, Skjaerven LH, Guitard Sein-Echaluce L, Catalan-Matamoros D. Effectiveness of movement and body awareness therapies in patients with fibromyalgia: a systematic review and meta-analysis. *Eur J Phys Rehabil Med.* 2019; 55(5):646–657. https://doi.org/10.23736/S1973-9087.19.05291-2.
- Busch AJ, Barber KAR, Overend TJ, Peloso PMJ, Schachter CL. Exercise for treating fibromyalgia syndrome. *Cochrane Database Syst Rev.* 2007. https://doi.org/ 10.1002/14651858.CD003786.PUB2/MEDIA/CDSR/CD003786/IMAGE\_N/ NCD003786-CMP-002-05.PNG. Published online.
- Hutton B, Salanti G, Caldwell DM, et al. The PRISMA Extension Statement for reporting of systematic reviews incorporating network meta-analyses of Health care interventions: checklist and explanations. *Ann Intern Med.* 2015;162(11): 777–784. https://doi.org/10.7326/M14-2385.
- Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol*. 2005;5(1):1–10. https://doi.org/10.1186/1471-2288-5-13/TABLES/3.

- 26. Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol*. 2014;14(1):1–13. https://doi.org/10.1186/1471-2288-14-135/ TABLES/3.
- Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther.* 2003;83 (8):713–721. https://doi.org/10.1093/PTJ/83.8.713.
- Ghai S, Ghai I, Schmitz G, Effenberg AO. Effect of rhythmic auditory cueing on parkinsonian gait: a systematic review and meta-analysis. *Sci Rep.* 2018;8(1):1–19. https://doi.org/10.1038/s41598-017-16232-5.
- Papakonstantinou T, Nikolakopoulou A, Higgins JPT, Egger M, Salanti G. CINeMA: software for semiautomated assessment of the confidence in the results of network meta-analysis. *Campbell Syst Rev.* 2020;16(1):e1080. https://doi.org/10.1002/ CL2.1080.
- Dworkin R, Turk D, Wyrwich K, et al. Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. *J Pain*. 2008;9(2):105–121.
- Albright J, Allman R, Bonfiglio R, et al. Philadelphia Panel evidence-based clinical practice Guidelines on selected rehabilitation interventions: overview and methodology. *Phys Ther.* 2001;81(10):1629–1640.
- Deeks J, Higgins J. Statistical algorithms in review manager 5. Stat Algorithm Rev Manager 5. 2007. Published online.
- Harrer M, Cuijpers P, Furukawa TA, Ebert DD. Doing meta-analysis with R : a hands-on guide. *Doing Meta-Anal R*. 2021. https://doi.org/10.1201/ 9781003107347. Published online.
- Salanti G, Ades AE, Ioannidis JPA. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. J Clin Epidemiol. 2011;64(2):163–171. https://doi.org/10.1016/J. JCLINEPI.2010.03.016.
- Rücker G, Schwarzer G. Ranking treatments in frequentist network meta-analysis works without resampling methods. *BMC Med Res Methodol*. 2015;15(1):58. https://doi.org/10.1186/s12874-015-0060-8.
- **36.** Shim S, Lee J. Network meta-analysis: application and practice using R software. *Epidemiol Health.* 2019:41.
- Hong H, Carlin BP, Shamliyan TA, et al. Comparing bayesian and frequentist approaches for multiple outcome mixed treatment comparisons. *Med Decis Making*. 2013;33(5):702–714. https://doi.org/10.1177/0272989×13481110.
- Rücker G, Krahn U, König J, Efthimiou O, Schwarzer G. Netmeta: network metaanalysis using frequentist methods. *R Package Version*. 2019;1(2).
- Viechtbauer W. Accounting for heterogeneity via random-effects models and moderator analyses in meta-analysis. J Psychol. 2007;215(2):104–121. https://doi. org/10.1027/0044-3409.215.2.104.
- Higgins J, Thompson S, Deeks J, Altman D. Measuring inconsistency in metaanalyses. *BMJ*. 2003;327(7414):557–560. https://doi.org/10.1136/ bmj.327.7414.557.
- Ter Veer E, van Oijen MGH, van Laarhoven HWM. The use of (Network) metaanalysis in clinical oncology. *Front Oncol.* 2019;9(AUG):822. https://doi.org/ 10.3389/FONC.2019.00822/BIBTEX.
- Mengshoel AM, Komnaes HB, Førre O. The effects of 20 weeks of physical fitness training in female patients with fibromyalgia. *Clin Exp Rheumatol.* 1992;10(4): 345–349. http://www.ncbi.nlm.nih.gov/pubmed/1395219.
- Häkkinen A, Häkkinen K, Hannonen P, Alen M. Strength training induced adaptations in neuromuscular function of premenopausal women with fibromyalgia: comparison with healthy women. *Ann Rheum Dis.* 2001;60(1):21–26. https://doi.org/10.1136/ard.60.1.21.
- Jentoft ES, Kvalvik AG, Mengshoel AM. Effects of pool-based and land-based aerobic exercise on women with fibromyalgia/chronic widespread muscle pain. *Arthritis Rheum.* 2001;45(1):42–47. https://doi.org/10.1002/1529-0131(200102) 45:1<42. :AID–ANR82>3.0.CO;2-A.
- Jones KD, Burckhardt CS, Clark SR, Bennett RM, Potempa KM. A randomized controlled trial of muscle strengthening versus flexibility training in fibromyalgia. *J Rheumatol.* 2002;29(5):1041–1048. http://www.ncbi.nlm.nih.gov/pubmed /12022321.
- 46. van Santen M, Bolwijn P, Landewé R, et al. High or low intensity aerobic fitness training in fibromyalgia: does it matter? *J Rheumatol*. 2002;29(3):582–587. http ://www.ncbi.nlm.nih.gov/pubmed/11908577.
- Schachter CL, Busch AJ, Peloso PM, Sheppard MS. Effects of short versus long bouts of aerobic exercise in sedentary women with fibromyalgia: a randomized controlled trial. *Phys Ther.* 2003;83(4):340–358. http://www.ncbi.nlm.nih.gov/p ubmed/12665405.
- Valim V, Oliveira L, Suda A, et al. Aerobic fitness effects in fibromyalgia. J Rheumatol. 2003;30(5):1060–1069. http://www.ncbi.nlm.nih.gov/pubmed /12734907.
- Sencan S, Ak S, Karan A, Muslumanoglu L, Ozcan E, Berker E. A study to compare the therapeutic efficacy of aerobic exercise and paroxetine in fibromyalgia syndrome. J Back Musculoskelet Rehabil. 2004;17(2):57–61. https://doi.org/ 10.3233/BMR-2004-17204.
- Valkeinen H. Changes in knee extension and flexion force, EMG and functional capacity during strength training in older females with fibromyalgia and healthy controls. *Rheumatology (Oxford)*. 2003;43(2):225–228. https://doi.org/10.1093/ rheumatology/keh027.
- Gusi N, Tomas-Carus P, Häkkinen A, Häkkinen K, Ortega-Alonso A. Exercise in waist-high warm water decreases pain and improves health-related quality of life and strength in the lower extremities in women with fibromyalgia. *Arthritis Rheum.* 2006;55(1):66–73. https://doi.org/10.1002/art.21718.

- Munguía-Izquierdo D, Legaz-Arrese A. Exercise in warm water decreases pain and improves cognitive function in middle-aged women with fibromyalgia. *Clin Exp Rheumatol.* 2007;25(6):823–830. http://www.ncbi.nlm.nih.gov/pubmed /18173915.
- Rooks DS, Gautam S, Romeling M, et al. Group exercise, education, and combination self-management in women with fibromyalgia: a randomized trial. *Arch Intern Med.* 2007;167(20):2192–2200. https://doi.org/10.1001/ archinte.167.20.2192.
- 54. Tomás-Carús P, Gusi N, Leal A, García Y, Ortega-Alonso A. The fibromyalgia treatment with physical exercise in warm water reduces the impact of the disease on female patients' Physical and mental health. *Reumatología Clínica (English Edition)*. 2007;3(1):33–37. https://doi.org/10.1016/S2173-5743(07)70206-X.
- Alentorn-Geli E, Padilla J, Moras G, Haro CL, Fernández-Solà J. Six weeks of whole-body vibration exercise improves pain and fatigue in women with fibromyalgia. J Alternat Complement Med. 2008;14(8):975–981. https://doi.org/ 10.1089/acm.2008.0050.
- Bircan Ç, Karasel SA, Akgün B, El Ö, Alper S. Effects of muscle strengthening versus aerobic exercise program in fibromyalgia. *Rheumatol Int.* 2008;28(6):527–532. https://doi.org/10.1007/s00296-007-0484-5.
- Bressan L, Matsutani L, Assumpção A, Marques A, Cabral C. Effects of muscle stretching and physical conditioning as physical therapy treatment for patients with fibromyalgia. *Revista Brasileira de Fisioterapia*. 2008;12(2):88–93. https://doi. org/10.1590/S1413-35552008000200003.
- Evcik D, Yigit I, Pusak H, Kavuncu V. Effectiveness of aquatic therapy in the treatment of fibromyalgia syndrome: a randomized controlled open study. *Rheumatol Int.* 2008;28(9):885–890. https://doi.org/10.1007/s00296-008-0538-3.
- Ide M, Laurindo I, Rodrigues-Junior A, Tanaka C. Effect of aquatic respiratory exercise-based program in patients with fibromyalgia. *Int J Rheum Dis.* 2008;11(2): 131–140. https://doi.org/10.1111/j.1756-185X.2008.00348.x.
- Tomas-Carus P, Gusi N, Häkkinen A, Häkkinen K, Leal A, Ortega-Alonso A. Eight months of physical training in warm water improves physical and mental health in women with fibromyalgia: a randomized controlled trial. J Rehabil Med. 2008;40 (4):248–252. https://doi.org/10.2340/16501977-0168.
- 61. Valkeinen H, Alén M, Häkkinen A, Hannonen P, Kukkonen-Harjula K, Häkkinen K. Effects of concurrent strength and endurance training on physical fitness and symptoms in postmenopausal women with fibromyalgia: a randomized controlled trial. Arch Phys Med Rehabil. 2008;89(9):1660–1666. https://doi.org/10.1016/j. apmr.2008.01.022.
- Altan L, Korkmaz N, Bingol Ü, Gunay B. Effect of pilates training on people with fibromyalgia syndrome: a pilot study. Arch Phys Med Rehabil. 2009;90(12): 1983–1988. https://doi.org/10.1016/j.apmr.2009.06.021.
- Carson JW, Carson KM, Jones KD, Bennett RM, Wright CL, Mist SD. A pilot randomized controlled trial of the Yoga of Awareness program in the management of fibromyalgia. *Pain.* 2010;151(2):530–539. https://doi.org/10.1016/j. pain.2010.08.020.
- 64. Kayo AH, Peccin MS, Sanches CM, Trevisani VFM. Effectiveness of physical activity in reducing pain in patients with fibromyalgia: a blinded randomized clinical trial. *Rheumatol Int.* 2011;32(8):2285–2292. https://doi.org/10.1007/s00296-011-1958-z.
- Baptista AS, Villela AL, Jones A, Natour J. Effectiveness of dance in patients with fibromyalgia: a randomized, single-blind, controlled study. *Clin Exp Rheumatol*. 2012;30(6):18–23. Suppl 74 http://www.ncbi.nlm.nih.gov/pubmed/23020850.
  López-Rodríguez M del M, Castro-Sánchez AM, Fernández-Martínez M, Matarán-
- 66. López-Rodríguez M del M, Castro-Sánchez AM, Fernández-Martínez M, Matarán-Peñarrocha GA, Rodríguez-Ferrer ME. Comparison between aquatic-biodanza and stretching for improving quality of life and pain in patients with fibromyalgia. *Aten Primaria*. 2012;44(11):641–649. https://doi.org/10.1016/j.aprim.2012.03.002.
- Letieri RV, Furtado GE, Letieri M, et al. Pain, quality of life, self perception of health and depression in patients with fi bromyalgia, submitted to hydrocinesiotherapy. *Rev Bras Reumatol.* 2013;53(6):494–500. https://doi.org/ 10.1016/j.rbr.2013.04.004.
- López-Rodríguez MM, Fernández-Martínez M, GA Matarán-Peñarrocha, et al. Effectiveness of aquatic biodance on sleep quality, anxiety and other symptoms in patients with fibromyalgia. *Med Clin (Barc)*. 2013;141(11):471–478. https://doi. org/10.1016/j.medcli.2012.09.036.
- Duruturk N, Tuzun EH. Culhaoglu B. Is balance exercise training as effective as aerobic exercise training in fibromyalgia syndrome? *Rheumatol Int.* 2015;35(5): 845–854. https://doi.org/10.1007/s00296-014-3159-z.
- Gavi MBRO, Vassalo DV, Amaral FT, et al. Strengthening exercises improve symptoms and Quality of life but do not change autonomic modulation in fibromyalgia: a randomized clinical trial. Cordero MD, ed. *PLoS One*. 2014;9(3), e90767. https://doi.org/10.1371/journal.pone.0090767.
- Genc A, Tur BS, Aytur YK, Oztuna D, Erdogan MF. Does aerobic exercise affect the hypothalamic-pituitary-adrenal hormonal response in patients with fibromyalgia syndrome? J Phys Ther Sci. 2015;27(7):2225–2231. https://doi.org/10.1589/ jpts.27.2225.
- Larsson A, Palstam A, Löfgren M, et al. Resistance exercise improves muscle strength, health status and pain intensity in fibromyalgia-a randomized controlled trial. Arthritis Res Ther. 2015;17(1). https://doi.org/10.1186/S13075-015-0679-1.
- 73. PÁ LatorreRomán, MA SantosE Campos, García-Pinillos F. Effects of functional training on pain, leg strength, and balance in women with fibromyalgia. *Mod Rheumatol.* 2015;25(6):943–947. https://doi.org/10.3109/ 14397595.2015.1040614.
- 74. Sañudo B, Carrasco L, de Hoyo M, Figueroa A, Saxton JM. Vagal modulation and symptomatology following a 6-month aerobic exercise program for women with fibromyalgia. *Clin Exp Rheumatol.* 2015;33(1). Suppl 88S41-5 http://www.ncbi. nlm.nih.gov/pubmed/25786042.

- Sevimli D, Kozanoglu E, Guzel R, Doganay A. The effects of aquatic, isometric strength-stretching and aerobic exercise on physical and psychological parameters of female patients with fibromyalgia syndrome. *J Phys Ther Sci.* 2015;27(6): 1781–1786. https://doi.org/10.1589/jpts.27.1781.
- Ekici G, Unal E, Akbayrak T, Vardar-Yagli N, Yakut Y, Karabulut E. Effects of active/passive interventions on pain, anxiety, and quality of life in women with fibromyalgia: randomized controlled pilot trial. *Women Health*. 2016;57(1): 88–107. https://doi.org/10.1080/03630242.2016.1153017.
- Fernandes G, Jennings F, Nery Cabral MV, Pirozzi Buosi AL, Natour J. Swimming improves pain and functional capacity of patients with fibromyalgia: a randomized controlled trial. Arch Phys Med Rehabil. 2016;97(8):1269–1275. https://doi.org/ 10.1016/j.apmr.2016.01.026.
- Alev A, Mihriban A, Bilge E, et al. Effects of whole body vibration therapy in pain, function and depression of the patients with fibromyalgia. *Complement Ther Clin Pract.* 2017;28:200–203. https://doi.org/10.1016/j.ctcp.2017.06.008.
- Assumpção A, Matsutani LA, Yuan SL, et al. Muscle stretching exercises and resistance training in fibromyalgia: which is better? A three-arm randomized controlled trial. *Eur J Phys Rehabil Med.* 2017;54(5):663–670. https://doi.org/ 10.23736/S1973-9087.17.04876-6.
- Collado-Mateo D, Dominguez-Muñoz FJ, Adsuar JC, Garcia-Gordillo MA, Gusi N. Effects of exergames on quality of life, pain, and disease effect in women with fibromyalgia: a randomized controlled trial. *Arch Phys Med Rehabil.* 2017;98(9): 1725–1731. https://doi.org/10.1016/j.apmr.2017.02.011.
- Toprak Celenay S, Anaforoglu Kulunkoglu B, Yasa ME, et al. A comparison of the effects of exercises plus connective tissue massage to exercises alone in women with fibromyalgia syndrome: a randomized controlled trial. *Rheumatol Int.* 2017;37 (11):1799–1806. https://doi.org/10.1007/s00296-017-3805-3.
- Acosta-Gallego A, Ruiz-Montero PJ, Castillo-Rodríguez A. Land- and pool-based intervention in female fibromyalgia patients: a randomized-controlled trial. *Turk J Phys Med Rehabil.* 2018;64(4):337–343. https://doi.org/10.5606/tftrd.2018.2314.
- Ernberg M, Christidis N, Ghafouri B, et al. Plasma cytokine levels in fibromyalgia and their response to 15 weeks of progressive resistance exercise or relaxation therapy. *Mediators Inflamm.* 2018. https://doi.org/10.1155/2018/3985154, 2018.
- Wong A, Figueroa A, Sanchez-Gonzalez MA, Son WM, Chernykh O, Park SY. Effectiveness of Tai Chi on cardiac autonomic function and symptomatology in women with fibromyalgia: a randomized controlled trial. J Aging Phys Act. 2018;26 (2):214–221. https://doi.org/10.1123/JAPA.2017-0038.
- Andrade CP, Zamunér AR, Forti M, Tamburús NY, Silva E. Effects of aquatic training and detraining on women with fibromyalgia: controlled randomized clinical trial. *Eur J Phys Rehabil Med.* 2019;55(1):79–88. https://doi.org/ 10.23736/S1973-9087.18.05041-4.
- Gomez-Hernandez M, Gallego-Izquierdo T, Martinez-Merinero P, et al. Benefits of adding stretching to a moderate-intensity aerobic exercise programme in women with fibromyalgia: a randomized controlled trial. *Clin Rehabil.* 2019;34(2): 242–251. https://doi.org/10.1177/0269215519893107.
- Jablochkova A, Bäckryd E, Kosek E, et al. Unaltered low nerve growth factor and high brain-derived neurotrophic factor levels in plasma from patients with fibromyalgia after a 15-week progressive resistance exercise. *J Rehabil Med.* 2019; 51(10):779-787. https://doi.org/10.2340/16501977-2593.
  Silva HJ de A, Assunção Júnior JC, de Oliveira FS, Oliveira JM de P, Figueiredo
- Silva HJ de A, Assunção Júnior JC, de Oliveira FS, Oliveira JM de P, Figueiredo Dantas G. Sophrology versus resistance training for treatment of women with fibromyalgia: a randomized controlled trial. *J Bodyw Mov Ther.* 2019;23(2): 382–389. https://doi.org/10.1016/j.jbmt.2018.02.005.
  Celenay ST, Mete O, Akan S, et al. Comparison of the effects of stabilization
- Celenay ST, Mete O, Akan S, et al. Comparison of the effects of stabilization exercise plus kinesio taping and stabilization exercise alone on pain and well-being in fibromyalgia. *Complement Ther Clin Pract.* 2020;38, 101076. https://doi.org/ 10.1016/j.ctcp.2019.101076.
- Villafaina S, Collado-Mateo D, Domínguez-Muñoz F, Fuentes-García JP, Gusi N. Benefits of 24-week exergame intervention on health-related quality of life and pain in women with Fibromyalgia: a single-blind, randomized controlled trial. *Games Health J.* 2019;8(6):380–386. https://doi.org/10.1089/g4h.2019.0023.
  Atan T, Karavelioğlu Y. Effectiveness of high-intensity interval training vs
- Atan T, Karavelioğlu Y. Effectiveness of high-intensity interval training vs moderate-intensity continuous training in patients with fibromyalgia: a pilot randomized controlled trial. Arch Phys Med Rehabil. 2020;101(11):1865–1876. https://doi.org/10.1016/j.apmr.2020.05.022.
- Britto A, Rodrigues V, dos Santos AM, et al. Effects of water- and land-based exercises on quality of life and physical aspects in women with fibromyalgia: a randomized clinical trial. *Musculoskeletal Care*. 2020;18(4):459–466. https://doi. org/10.1002/msc.1481.
- 93. de Medeiros SA, de Almeida, Silva HJ, do Nascimento RM, da Silva, Maia JB, de Almeida Lins CA, de Souza MC. Mat Pilates is as effective as aquatic aerobic exercise in treating women with fibromyalgia: a clinical, randomized and blind trial. Adv Rheumatol. 2020;60(1):21. https://doi.org/10.1186/s42358-020-0124-2.
- Gulsen C, Soke F, Eldemir K, et al. Effect of fully immersive virtual reality treatment combined with exercise in fibromyalgia patients: a randomized controlled trial. Assist Technol. 2020. https://doi.org/10.1080/ 10400435.2020.1772900. Published online.
- Sarmento CV, Moon S, Pfeifer T, et al. The therapeutic efficacy of Qigong exercise on the main symptoms of fibromyalgia: a pilot randomized clinical trial. *Integr Med Res.* 2020;9(4). https://doi.org/10.1016/j.imr.2020.100416.
- de CarvalhoMS, LC Carvalho, Menezes FDS, Frazin A, Gomes EDC, Iunes DH. Effects of exergames in women with fibromyalgia: a randomized controlled study. *Games Health J.* 2020;9(5):358–367. https://doi.org/10.1089/g4h.2019.0108.
- Swar W. Effect of underwater exercises on treating postmenopausal fibromyalgia symptoms. *Eur J Mol Clin Med.* 2020;7(9):661–672, 02251128.

- Arakaki J, Jennings F, Estrela GQ, Cruz Martinelli V d G, Natour J. Strengthening exercises using swiss ball improve pain, health status, quality of life and muscle strength in patients with fibromyalgia: a randomized controlled trial. *Reumatismo*. 2021;73(1):15–23. https://doi.org/10.4081/reumatismo.2021.1357.
- Garrido-Ardila EM, González-López-arza MV, Jiménez-Palomares M, García-Nogales A, Rodríguez-Mansilla J. Effects of physiotherapy vs. Acupuncture in quality of life, pain, stiffness, difficulty to work and depression of women with fibromyalgia: a randomized controlled trial. J Clin Med. 2021;10(17). https://doi. org/10.3390/jcm10173765.
- Hernando-Garijo I, Ceballos-Laita L, Mingo-Gomez M, et al. Immediate effects of a telerehabilitation program based on aerobic exercise in women with fibromyalgia. *Int J Environ Res Public Health*. 2021;18(4). https://doi.org/10.3390/ ijerph18042075.
- 101. Rodríguez-Mansilla J, Mejías-Gil A, Garrido-Ardila EM, et al. Effects of nonpharmacological treatment on pain, flexibility, balance and quality of life in women with fibromyalgia: a randomised clinical trial. *J Clin Med.* 2021;10(17). https://doi.org/10.3390/jcm10173826.
- 102. Schachter CL, Busch AJ, Peloso PM, et al. Effects of short versus long bouts of aerobic exercise in sedentary women with fibromyalgia: a randomized controlled trial. *Phys Ther.* 2003;83(4):340–358.
- 103. Atan T, Karavelioğlu Y, Atan T, Karavelioğlu Y. Effectiveness of high-intensity interval training vs moderate-intensity continuous training in patients with fibromyalgia: a pilot randomized controlled trial. Arch Phys Med Rehabil. 2020;101 (11):1865–1876. https://doi.org/10.1016/J.APMR.2020.05.022.
- 104. Rooks DS, Gautam S, Romeling M, et al. Group exercise, education, and combination self-management in women with fibromyalgia: a randomized trial. *Arch Intern Med.* 2007;167(20):2192–2200. https://doi.org/10.1001/ ARCHINTE.167.20.2192.
- Alentorn-Geli E, Padilla J, Moras G, Haro CL, Fernández-Solà J. Six weeks of wholebody vibration exercise improves pain and fatigue in women with fibromyalgia.

J Altern Complement Med. 2008;14(8):975–981. https://doi.org/10.1089/ ACM.2008.0050.

- Lin L, Zhang J, Hodges JS, Chu H. Performing arm-based network meta-analysis in R with the pcnetmeta package. J Stat Softw. 2017;80. https://doi.org/10.18637/ JSS.V080.105.
- 107. Fink L, Lewis D. Exercise as a treatment for fibromyalgia: a scoping review. J Nurse Pr. 2017;13(8):546–551.
- 108. Vaegter H, Jones M. Exercise-induced hypoalgesia after acute and regular exercise: experimental and clinical manifestations and possible mechanisms in individuals with and without pain. *Pain Rep.* 2020;5(5):e823.
- Newcomb L, Koltyn K, Cook D. The influence of a preferred versus a prescribed exercise session on pain in women with fibromyalgia. J Pain. 2009;10(4):S16. https://doi.org/10.1016/j.jpain.2009.01.070.
- Arakaki J, Jennings F, Estrela G, Martinelli V, Natour J. Strengthening exercises using swiss ball improve pain, health status, quality of life and muscle strength in patients with fibromyalgia: a randomized controlled trial. *Reumatismo*. 2021;73(1): 15–23. https://doi.org/10.4081/REUMATISMO.2021.1357.
- 111. Garrido-Ardila EM, González-López-arza MV, Jiménez-Palomares M, García-Nogales A, Rodríguez-Mansilla J. Effects of physiotherapy vs. Acupuncture in quality of life, pain, stiffness, difficulty to work and depression of women with fibromyalgia: a randomized controlled trial. J Clin Med. 2021;10(17). https://doi. org/10.3390/jcm10173765.
- 112. Hernando-Garijo I, Ceballos-Laita L, Mingo-Gomez M, et al. Immediate effects of a telerehabilitation program based on aerobic exercise in women with fibromyalgia. *Int J Environ Res Public Health*. 2021;18(4). https://doi.org/10.3390/ ijerph18042075.
- 113. Pontes-Silva A. Fibromyalgia: a new set of diagnostic criteria based on the biopsychosocial model. *Rheumatology (Oxford)*. 2024. https://doi.org/10.1093/ rheumatology/keae074. Published online February 3.