Contents lists available at ScienceDirect



Masterclass

Brazilian Journal of Physical Therapy



journal homepage: www.elsevier.com/locate/bjpt

Rehabilitation management of the chronic pain-hypertension synergy: Proposal of an evidence-informed framework

Renzo Mendoza^{a,*}^o, Craig Hensley^b, Jennifer Ryan^b, Todd Davenport^c

^a Department of Therapy Services, University of Illinois Health Systems, 711 W. Maxwell St. Chicago, IL 60607 USA

^b Department of Physical Therapy and Human Movement Sciences, Northwestern University, 645 N. Michigan Ave., Suite 1100 Chicago, IL 60611 USA

^c School of Health Sciences, University of the Pacific, Stockton, California 3601 Pacific Ave., Stockton, CA 95211 USA

ARTICLE INFO

Keywords: Cardiovascular disease Epidemiology Examination Exercise Physical therapy Risk factors

ABSTRACT

Background: Chronic pain and hypertension (HTN) have a significant societal impact worldwide. Both lead to consumption of substantial healthcare resources and dollars. The pathophysiology of chronic pain and HTN are each complex and multifactorial. Chronic pain has been associated with an increased risk of comorbid HTN. There are numerous common factors related to the development, progression, and prognosis of patients with chronic pain and HTN. Though the cardiovascular and pain regulatory systems are structurally and functionally intertwined, the exact physiological interactions between chronic pain and HTN are less understood. Many systems, including the central and autonomic nervous systems, play a role in the development and interaction of these two complex conditions. In collaboration with other healthcare providers, physical therapists play a critical role in managing those with chronic pain and HTN.

Objectives: This masterclass aims to develop physical therapists' knowledge and understanding regarding 1) the epidemiological and physiological relationship between chronic pain and HTN, and 2) screening, examination, and management of the patient with chronic pain and HTN, including appropriate referral.

Discussion: Improving understanding of the relationship between chronic pain and HTN should assist clinicians in appropriate management strategies for such patients. Physical therapy should be an essential component in the management of patients with chronic pain and HTN. By collaboratively addressing both health conditions, there is a window of opportunity to play a role in mitigating the burden of these conditions and promoting a healthier society.

Introduction

Chronic pain and hypertension (HTN) are worldwide epidemics that can be challenging to manage. Chronic pain affects 20 % of people, while HTN affects nearly one-third of the global adult population.^{1,2} Chronic pain has been associated with multiple physical and mental health conditions, lost productivity, higher suicide rate, and substance abuse.³⁻⁷ HTN prevalence has increased across the world since its reclassification. The updated American Heart Association and American College of Cardiology defines HTN as a systolic blood pressure (BP) \geq 130mmHg or a diastolic BP \geq 80mm Hg. Though the exact etiology of elevated blood pressure and HTN is often not known, the effectiveness of risk reduction through lifestyle and dietary changes make it the most common preventable non-communicable disease that leads to cardiovascular disease and all-cause mortality.³⁻⁷ The total costs of chronic pain and HTN are substantial across the globe. Healthcare expenditures in the United States alone range from \$261-300 billion and \$131 billion for chronic pain and HTN, respectively.^{8,9}

Physical therapists can play a pivotal role in managing patients with chronic pain and HTN. In this master class, we discuss the current literature regarding the relationship between chronic pain and HTN at the epidemiological and pathophysiological levels. In addition, we provide evidence suggesting that the two health conditions together comprise a syndemic.¹⁰ Further, we propose evidence-based guidelines regarding the screening, examination, and non-pharmacological management of patients with comorbid chronic pain and HTN.

Received 20 July 2024; Received in revised form 20 December 2024; Accepted 14 May 2025 Available online 3 June 2025

^{*} Corresponding author at: 711 W. Maxwell St. Chicago, IL 60607 USA.

E-mail addresses: Renzomendozadpt@gmail.com (R. Mendoza), craig.hensley1@northwestern.edu (C. Hensley), jennifer-ryan@northwestern.edu (J. Ryan), tdavenport@pacific.edu (T. Davenport).

https://doi.org/10.1016/j.bjpt.2025.101229

^{1413-3555/© 2025} The Authors. Published by Elsevier España, S.L.U. on behalf of Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

The relationship between chronic pain and hypertension at the population level

Globally, up to 4 in 10 people with chronic pain have HTN.¹¹ Relationships have been found between HTN and osteoarthritis, chronic low back pain (LBP), rheumatoid arthritis (RA), fibromyalgia, and complex regional pain syndrome.^{2,11} Chronic pain also has a bidirectional relationship with several hypertensive risk factors, including metabolic syndrome, suboptimal diet, poor sleep quality, elevated body mass index (BMI), abdominal obesity, Type 2 diabetes, cigarette smoking, social determinants of health, and mental health conditions.^{2,12,13} The relationship between chronic pain, HTN, and associated factors is illustrated in Fig. 1.

Individuals with chronic pain and cardiovascular disease report greater pain intensity and disability compared to those suffering from chronic pain alone.^{2,14} Moreover, a recent cross-sectional study has shown that the presence of comorbidities is associated with higher odds of daily pain, pain impact, and pain intensity.¹⁵ Individuals with chronic pain and comorbidities are less likely to receive appropriate care, have a poorer prognosis, and account for higher payment cost.^{16,17} Further, many of the associated factors listed in Fig. 1 have been shown to influence individuals' pain intensity, symptom distribution, and pain reactivity.^{2,15}

The interplay between chronic pain and HTN can be considered a syndemic. This term refers to the coexistence and interaction of multiple health conditions and factors that synergistically worsen prognosis and outcomes, creating a compounded health burden greater than the sum of its parts.¹⁰ Understanding this relationship allows physical therapists to develop more holistic and effective interventions that address these

intertwined factors, ultimately leading to better patient outcomes.

Potential mechanisms mediating the relationship between blood pressure and pain

Hemodynamic response to noxious stimuli in healthy individuals

The cardiovascular and pain regulatory systems are structurally and functionally intertwined. In healthy individuals, these interconnections produce blood pressure (BP)-related hypoalgesia, in which a noxious stimulus triggers elevated BP to initiate a reduction in pain experience (Supplemental material 1).^{18,19} A noxious stimulus travels through the somato-autonomic reflex, which increases BP through sympathetic arousal (Fig. 2a). Increased BP leads to baroreceptor stimulation, triggering the baroreflex and associated pathway in the central nervous system (CNS). The baroreceptor system is vital to cardiovascular regulation. It consists of a negative feedback loop in which electrical activity changes in arterial baroreceptors, resulting from BP alteration, initiate compensatory changes in heart rate, cardiac contractility, and vascular tone.²⁰ This feedback system processes information within the cardiovascular cranial nerve branches into the lower brain stem. Brain signaling reinforces sympathetic output and descending modulation of noxious stimuli via noradrenergic pathways to reduce the painful experience. Once reduction in noxious and pain signaling occurs, there is a reduction in activation within this baroreflex and associated pathways, allowing BP to return to baseline.

Although BP-related hypoalgesia has been postulated to respond to acute pain stimulus, it is important to highlight this mechanism has demonstrated mixed results in conjunction with exercise. Exercise



Fig. 1. The relationship between chronic pain, hypertension, and associated factors.



Fig. 2a. (A) Along with the secondary projections that sends signals to supraspinal centers, the noxious stimulus additionally travels through the somato-autonomic reflex, which increases sympathetic arousal and produces increased blood pressure. (B) Increased blood pressure leads to baroreceptor stimulation which triggers the baroreflex, inputting sensory information into the central nervous system. Alterations in baroreceptor sensitivity may limit the ability for the autonomic system to stimulate the central autonomic network which includes the nucleus of the solitary tract and the locus coeruleus. (C) Signals from the baroreceptor triggers the central autonomic network. (D) This reinforces sympathetic output and pain inhibition via noradrenergic pathways, decreasing pain stimulus. Adapted with permission from Bruehl et al. pain and blood pressure model²⁶.



Fig. 2b. Alterations in the pain and blood pressure model may include noradrenergic exhaustion where a reduction in norepinephrine which normally modulates pain. Another possibility in this region includes facilitatory mechanisms with changes in receptor type may promote symptoms. Adapted with permission from Bruehl et al. alterations in the pain and blood pressure model²⁶.

inherently is thought to produce an increase in BP that attenuates the baroreceptor related mechanism.²¹ Research on isometric exercises found that increases in BP has not consistently corresponded to alterations in pain perception.²² These findings suggest that other physiological mechanisms may be at play aside BP alone such as endogenous opioids, cannabinoids and growth hormones which have been hypothesized to be produced during exercise.²¹

Alterations in hemodynamic response to noxious stimuli in individuals with chronic pain

Altered central nociceptive processing may occur with chronic pain, leading to nociplastic pain.²³ While various mechanisms are associated with nociplastic pain, including hyperexcitability of nociceptive pathways, another potential feature of the condition is impaired descending inhibitory mechanisms. Descending noradrenergic inhibition can be profoundly effective in modulating pain, and may be impaired in individuals with HTN and chronic pain.^{24,25} Chronic pain is associated with reduced magnitude of BP-related hypoalgesia, suggesting maladaptive processes occurring within the cardiovascular regulatory systems (Supplemental Material 1). These are potentially relevant to comorbid cardiovascular risk in chronic pain.²⁶ Possible reasons for this altered relationship between BP and pain sensitivity include exhaustion of certain pain-inhibiting pathways, changes in baroreceptor function, increased inflammation, and other contributing factors (Fig. 2b).

Noradrenergic exhaustion

In chronic pain, persistent and excessive antinociceptive or endogenous pain blocking demands may eventually exhaust noradrenergic pathways²⁴⁻²⁶ (Fig. 2b). It additionally has been hypothesized that persistent nociceptive input and depletion of norepinephrine reserve leads to a reduction of receptor types within the dorsal horn. These two factors lead to an inability of norepinephrine to stimulate pain inhibition at the spinal level. Thus, pain is left uninhibited, allowing for a continuous cycle to increase BP. A reduction in noradrenergic receptors has also been shown to contribute to HTN by affecting renal function.²⁷ When the renin angiotensin aldosterone system is overly active, it can cause persistent vasoconstriction, increase blood volume, overstimulate the sympathetic nervous system, and cause vascular remodeling.^{28,29} This can lead to many systemic conditions, including essential HTN, renovascular HTN, heart failure, and chronic kidney disease.²⁹

Noradrenergic facilitation

Alterations in noradrenergic facilitatory mechanisms result in noradrenaline producing more nociceptive drive rather than inhibiting (Fig. 2b). This stems in part from descending facilitatory mechanisms and/or promotion of pain input.^{30,31} Persistent pain has been shown to increase excitability and sensitivity of spinal neurons, further driving nociplastic changes.³²⁻³⁵ These CNS changes also include alterations in receptor type which have been hypothesized to increase firing when bound by norepinephrine. Noradrenergic facilitation increases spinal cord excitability when exposed to an influx of norepinephrine. Pain is further facilitated, with continuous driving of BP elevation.

Baroreceptor habituation

It is important to highlight that baroreceptor efficiency is often determined through the measurement of baroreceptor sensitivity.³⁶ This measurement is defined as the change in the R-R interval in milliseconds per mmHg change in systolic BP and is the most established parameter describing the function of the baroreflex loop.³⁶ Reductions in baroreceptor sensitivity have been demonstrated in those with HTN³⁶ and chronic pain states, including LBP, RA, fibromyalgia, temporomandibular disorders, and irritable bowel syndrome.^{26,37} Reduced sensitivity has also been observed in the presence of several health conditions and other factors such as anxiety, diabetes, chronic smoking, and obesity.^{36,38} Alterations in pain regulatory processes associated with

chronic pain can be associated with changes in baroreceptor sensitivity.³⁶ Changes in baroreceptor sensitivity may occur due to an altered baroreceptor firing threshold. Thus, with changes in baroreceptor activation, an inability to reach the central nervous system occurs and ultimately influences pain inhibition (Fig. 2b).

Systemic inflammation

A bi-directional relationship is possible between systemic inflammation and chronic pain and/or HTN. Elevated levels of systemic inflammation have been associated with higher odds of having pain and contributes to initiation, progression, and maintenance of HTN.³⁹⁻⁴³ Chronic pain has been shown to contribute to a persistent inflammatory state at both the peripheral and central level. This results in neuroplastic changes, including peripheral and central sensitization.⁴⁴ HTN has also been proposed to trigger low grade systemic and vascular inflammation, leading to oxidative stress, and alterations in the extracellular matrix composition.^{45,46}

Other contributory factors

Stress, anxiety, sleep deprivation, and depression have been suggested to play a role in this mechanistic model linking chronic pain and HTN (Supplementary Material 2).¹⁸ These factors all relate to chronic stress and psychological distress which, both in animal and human studies, are related to CNS dysregulation and a neuroinflammatory state linked to chronic pain syndromes and HTN.^{37,47} This relationship underscores the importance of a comprehensive biopsychosocial screen, a practice that has long been an integral piece of the physical therapy examination.^{23,48,49}

Patient assessment

Evaluate for health conditions requiring referral

Chronic pain conditions may mask sinister pathology requiring assessment and intervention from another health care practitioner. Unfortunately, clinicians are more likely to discount symptoms that patients with chronic pain report and are less inclined to provide intervention, particularly when evidence for a particular pathology is lacking and/or the patient reports high pain intensity.^{50,51} It is imperative that health care providers continue to screen and examine throughout the episode of care, even if patients are improving. For instance, Hensley and Emerson report on a patient with chronic left upper quarter pain who was improving with management provided by the physical therapist. A change in presentation during the episode of care warranted referral for further medical workup, leading to a diagnosis of lung cancer.⁵² Moreover, patients with chronic pain and HTN should be evaluated for signs and symptoms consistent with post-exertional malaise/post-exertional neuroimmune exhaustion (PEM/PENE),⁵³ because pain is a common component of PEM/PENE. PEM/PENE is a clinical hallmark of myalgic encephalomyelitis (ME) and ME-like conditions, such as subtypes of Long Covid⁵⁴⁻⁵⁷ and post-treatment Lyme disease.⁵⁸ These conditions are important to be aware of because exercise recovery responses are impaired in people with PEM/PENE.⁵⁹⁻⁶³ In those with PEM/PENE, exercise should not be considered as a frontline intervention.⁶⁴

Identify predominant pain mechanism(s) involved

A mechanism-based approach to pain has been proposed to improve pain management. A mechanisms-based approach integrates and builds on the biopsychosocial models by defining specific pathobiology in pain processing, and external factors influence an individuals' pain experience.²³ Pathobiological categories in pain processing include nociceptive, nociplastic, and neuropathic pain. These are further defined in Table 1. Recent recommendations have proposed features that discriminate between these three categories.^{65,66} It is important to

Table 1

Identifying and differentiating between pain mechanisms^{65,70}.

	Nociceptive	Nociplastic	Neuropathic
Pain location	Local	Diffuse, widespread, multisite, and/or poorly localized. Varving distribution over time	Distributed to nerve injury/ pathology
Mechanism of injury	Clear, proportionate	Less clear or unclear, disproportionate	Consistent with nerve injury/ pathology
Pain behavior	Proportionate to injury and healing stage	Higher irritability Prolonged healing time	Consistent with nerve injury/ pathology
Pain descriptors	Sharp in acute stage Dull ache, throb at rest	Combination of nociceptive/ neuropathic descriptors	Lancinating, shooting, electric-like, burning, hot
Other subjective findings	+ response to anti-inflammatory medication	Fatigue, sleep, cognitive, mood, and/or gastrointestinal disturbance(s) Hypersensitivity to non-musculoskeletal stimuli (e.g., sound, heat, cold)	Paresthesia
Example screening tools		Central Sensitization Inventory score above ⁷⁰	 painDETECT⁷¹⁻⁷³ Leads Assessment of Neuropathic Signs and Symptoms⁷⁴
Movement testing	 Pain reported in particular phases of movement Consistent with a specific health condition 	 Pain reported through multiple phases of movement with multiple movement problems Less consistent or inconsistent with a specific health condition(s) 	Consistent with nerve injury/ pathology
Other objective findings	 Warm, swelling, skin color changes indicative of local inflammation Positive special tests linked to body structure(s) 	 Hyperalgesia and/or allodynia Pain pressure thresholds- lower at distant site, contralateral extremity, or throughout upper/lower quarter) Conditioned pain modulation- reduced or no change Temporal summation- increased 	 Positive central or peripheral neural exam findings Muscle performance Sensation Deep tendon reflexes Upper motor neuron testing Neurodynamics Other neural testing (i.e. tapping over

Nociceptive pain- pain arising from activation of nociceptors in non-neural tissues, which is traditionally associated with acute injury, inflammation, or mechanical irritant.

Nociplastic pain- pain that stems from alterations of nociceptive processing within the central nervous system and often there is no clear evidence of tissue damage nor somatosensory disease.

Neuropathic pain symptoms associated with a specific lesion or disease within the somatosensory system.

recognize that more than one category can simultaneously contribute to an individual's pain experience. In addition, consideration of pain-relevant psychological and motor factors must be given, as they may play a role in the maintenance of an individual's pain.²³ Interventions have been proposed that can be matched to specific pain mechanisms.^{23,67} For instance, a patient with LBP predominantly driven by a nociceptive source can be managed with interventions that target biomechanics, such as posture education, strengthening weak musculature, and stretching structures deemed inflexible/hypomobile. Alternatively, if the pain is driven predominantly from a nociplastic source, it is unlikely that treatment targeting mechanics alone will improve outcomes. Pain neuroscience education (PNE) can be a crucial addition to exercise and manual therapy to assist these patients.68,69 It should be noted that there is a paucity of evidence demonstrating that a mechanism-based approach to pain provides better benefits to patient outcomes compared to a thorough biopsychosocial assessment and person-centered management plan.

While the standard patient interview and physical examination is critical, physical therapists may use additional testing, including quantitative sensory testing (QST), to help determine pain mechanism(s) involved. These tests include palpation, pain pressure threshold, temporal summation, and conditioned pain modulation. It should be noted that these tests have several limitations as stand-alone pieces of an evaluation.^{12,75} Firstly, they are indirect measures of CNS excitability and inhibition of pain pathways. Secondly, there is a general lack of consensus on normative and predictive values. Thirdly, findings from these tests do not always correlate with symptoms intensity or improvement in symptoms.^{76–78} Finally, funding for equipment and time to perform QST may not be widely available across rehabilitation settings. There are, however, cheaper, alternative testing equipment that can be used to perform QST measures.^{79,80}

Palpation

Palpating the area where pain is reported and areas remote from the site of pain can provide valuable clues. With palpation, clinicians should be looking for hyperalgesia (i.e., augmented response to noxious stimuli) and allodynia (i.e., pain evoked with a non-noxious stimulus). If either of these palpation tests are positive, particularly distant from the site of pain, a nociplastic pain mechanism should be considered.^{23,80}

Pain pressure threshold (PPT)

PPT testing is a method used to objectively measure hyperalgesia. With the use of a pain algometer, pressure can be applied at sites local and distant to the pain location at a rate of about 30 kPa/s. Alternatively, a modified 10 ml syringe can be used as a substitute.⁸¹ Moderate evidence has shown that there may be a relationship between brain alteration and PPT.⁸² Further, patients are less likely to respond to physical therapy management when a reduction in PPT is found.⁸³

Conditioned pain modulation

The clinician measures PPT at a distant site before and after inducing a noxious stimulus (BP cuff or cold-water immersion). This is a proxy measure of descending pain inhibition and exercise-induced hypoalgesia (EIH). If PPT is reduced or no change is found, a nociplastic condition is suspected.⁸⁴

Temporal summation

Temporal summation is the term used to describe the phenomenon of windup. Windup is characterized by the gradual amplification activity in the dorsal horn cells following repetitive stimulation of primary afferent C-fibers.⁸¹ A repetitive noxious stimulus with a 0.7 mm hair/filament is applied with a frequency of 1/s for 30s in an area of hyperalgesia. Patients are asked to rate the pain intensity induced by the first and last stimuli.⁸¹ An increase in pain intensity between the first and last stimulus indicates enhanced central excitability and reflects facilitated

temporal summation. This response is considered a clinical sign of central sensitization. An alternative proxy measure could be a functional test, such as the six-minute walk test.⁸⁵ Elevation in temporal summation suggests that the patient is less likely to respond to physical therapy intervention.⁸³

Assessment of vital signs

Physical therapists possess the expertise to assess vital signs throughout the episode of care, not only to conduct health screenings and gather patient history but also to ensure safety and appropriateness prior to initiating exercise interventions. They are positioned to expedite the World Health Organization's initiative to reduce prevalence of undiagnosed or uncontrolled HTN.¹ The specific parameters to attend to in order to mitigate threats to accurate BP measurement is available from the American Heart Association for reference.⁸⁶ Appropriate BP management is likelv to reduce the risk of negative cardiovascular/cerebrovascular-related sequelae, and improve societal health.⁸⁷⁻⁹⁰ Vital sign assessment should include monitoring at rest, during aerobic activity, and during recovery. This testing approach can reveal BP readings that place the patient at risk for an adverse

event.^{1,12,16,91,92} Fig. 3 provides guidelines for assessment and management based on patient vitals during different stages of rest, aerobic testing/exercise, and during recovery to ensure safety and optimal health outcomes. It includes specific BP measures and corresponding clinical decision making.

Resting blood pressure

At rest, normal BP (<120/80 mmHg) allows for safe exercise testing if there is no history of organ disease or failure; any uncertainty requires consultation with another healthcare provider. Any reading between >120/80 and <180/120 mmHg in a patient without HTN diagnosis and symptoms (dizziness, nausea, tinnitus, headache, or pain unrelated to their primary diagnosis) necessitates at home BP measurements **for at least 1 week**, close in clinic monitoring, and referral to an appropriate healthcare provider for additional monitoring (usually 2-3 office visits at 1-4 week intervals).⁹³

In patients with known HTN, or elevated BP (120-129/<80 mmHg), at home and close in clinic monitoring is required throughout the episode of care. Readings between 130-159/80-99 mmHg in the patient with diagnosed HTN and on medication is considered to be uncontrolled. If asymptomatic, submaximal aerobic testing may proceed,



Fig. 3. Decision-making algorithm to identify and manage patients with elevated blood pressure.

education on lifestyle modification that is evidenced to reduce BP, and a referral to the healthcare practitioner managing the patients BP (general practitioner, cardiologist, etc) is required to ensure medical management is optimal. If a patient presents with symptoms such as headache, chest pain, tinnitus, or vision changes, exercise testing is contraindicated, and prompt medical evaluation is required. Any systolic reading >180 mmHg and/or diastolic reading >110 mmHg is a contraindication to resistance training and aerobic activity.94,95 The literature varies on the upper limit of BP readings as contraindications to exercise. However, we recommend the lowest limit as a conservative approach in individuals with chronic pain, as there are likely many other variables that impact exercise intensity in this population. Asymptomatic patients should seek care for management options within 1-2 days after detection, as this is considered a hypertensive urgency case. These patients could still benefit from other interventions prior to comprehensive testing. If the patient is symptomatic or shows signs of organ damage, this is considered a hypertensive emergency requiring access to emergency care.9

Blood pressure during aerobic activity

There is growing evidence for the utility of high BP responses to submaximal aerobic activity (60-70 % maximum HR intensity).98-100 Recent data have shown that systolic BP ≥160 mmHg during submaximal exercise testing is associated with an increased risk for a major cardiovascular event.^{98,101} During aerobic activity, normal vital responses include a gradual increase in systolic BP of approximately 10 mmHg per metabolic equivalent (MET) and a corresponding rise in heart rate (HR) of around 10 beats per minute (BPM) per MET as measured in normal adults.¹⁰² These increases reflect the body's appropriate physiological adaptation to the increased demand for oxygen during physical activity. Estimated MET levels using the Fitness Registry and the Importance of Exercise National Database (FRIEND) equations can assist the clinician with accurately estimating MET level increases.¹⁰³ For instance, walking at 2.41 kilometers per hour (km/h) at 0 % grade is equal to approximately 3 METs. Increasing speed to 3.22 km/h on a 3.5 % grade equates to approximately 4 METs. In patients taking beta blocker medication, HR response is blunted. The rate of perceived exertion (RPE) scale should be used in supplement to vital sign measurement.¹⁰⁴ The RPE is also a useful tool to guide exercise prescription in cases where the patient cannot tolerate aerobic testing. Regardless, an unexplained drop in systolic BP, no increase in systolic BP, or an exaggerated elevation in BP or HR may suggest poor cardiovascular response, deconditioning, or other potential cardiovascular abnormalities. The largest study available in the literature that measured exercise testing responses in patients with cardiovascular disease offers a sense of the variance from normal responses as driven by disease.¹⁰⁵ There is no database for standardized exercise testing responses for individuals with chronic pain. The clinician will need to allow self-selected intensities with ongoing vital sign monitoring due to variations in exercise intensity as driven by pain intensity.

If a patient without a history of HTN is found to have elevated BP in response to aerobic activity, careful monitoring of vitals over time is needed to confirm a diagnosis of HTN. In patients with HTN, abnormal responses require immediate cessation of aerobic activity with prompt referral for a medical evaluation to assess the underlying causes and determine if further diagnostic testing or treatment is needed. Additionally, if there are concerns about the patient's cardiovascular health, further monitoring and a more comprehensive evaluation may be warranted. Aerobic activity should be terminated if the patient develops symptoms (dyspnea, extreme fatigue, faintness, chest pain, unsteadiness, change in mentation), signs of poor peripheral perfusion (cyanosis, cramping, or pallor), there is a drop in systolic BP > 10mmHg, or there is a BP reading that exceeds 250/115 mmHg.106 Finally, it should be noted that manual measurement of BP is necessary during exercise. Automatic oscillometric BP measurement, while useful at rest, cannot accurately measure BP during exercise.¹⁰

Blood pressure during recovery

There is also growing utility for the use of BP and HR response after aerobic activity.^{87-90,98,108-110} For instance, in middle-aged men, if systolic BP is >195 mmHg two minutes after aerobic testing, there is a 69 % increased risk for a future myocardial infarction.¹¹¹ In contrast, recent data have shown a 1.27 increases odds of a major cardiovascular event for every 10 mmHg increase in diastolic BP three minutes after exercise.¹¹² This is understandable because an elevation in diastolic BP reduces the opportunity for the myocardial perfusion between beats. BP should be taken at completion of aerobic activity.¹¹³ BP should progressively trend to normal resting levels over the course of 5-6 min post-aerobic activity. HR should decrease at least 12 BPM in the first minute and return to near resting.¹¹³ If BP remains elevated beyond 5-6 min during recovery, it is considered an abnormal response and may signal an underlying issue, such as persistent HTN or poor cardiovascular conditioning. In such cases, the patient should undergo further medical evaluation.

Determine contributing factors to chronic pain and HTN

Multiple common risk factors should be considered when evaluating a patient with chronic pain and HTN. Assessment tools are identified and described in Table 1. As able, the health care team should work with the patient to mitigate these risk factors. Providers should also consider pain medications that can elevate BP. Risk of systolic/diastolic HTN can increase with consumption of non-steroidal anti-inflammatory drugs (e.g., naproxen), serotonin norepinephrine reuptake inhibitors (e.g., dulox-etine), and tricyclic antidepressants (e.g., amitriptyline).^{93,114}

Intervention for the patient with comorbid chronic pain and hypertension

Given the relationship described above between chronic pain and HTN, interventions could be designed to target risk factors for chronic pain and HTN and the health conditions themselves. Further, interventions targeting pain management may also reduce BP.

Education

Educational material should be provided in an accessible format to the patient. Education should be provided regarding:

- 1) What their BP reading means relative to healthy BP.
- 2) How to accurately measure BP at home for consistent comparison.3) Warning signs and symptoms that require a consultation with a physician or an emergency room visit.
- 4) The potential bi-directional relationship between chronic pain and HTN, and how management targeting a patient-centered approach can assist with management of both.
- 5) Associated risk factors, particularly modifiable risk factors like stress, sleep, physical activity, and diet, related to chronic pain and HTN. Stress, for example, can be addressed through teaching coping strategies using the eliminate-change-accept approach and/or by increasing greenspace exposure.¹¹⁵ Educating on the importance of sleep and sleep hygiene can be tailored to address any sleep-related dysfunction.¹¹⁶ Clinicians should consider a referral to a psychologist, sleep specialist, and/or dietitian when deemed appropriate.
- 6) Factors related to predominant pain mechanism(s). Modifiable biomechanical factors that may be further perpetuating symptoms should be discussed. PNE could be incorporated when the clinical picture is dominated by nociplastic pain, or when maladaptive pain perceptions and/or coping strategies are present.⁶⁸ Clinicians should aim to improve patients' knowledge of pain processing within the nervous system, engaging patients in a discussion regarding their pain and attempting to reconceptualize the patient's understanding and beliefs about their pain as needed.¹¹⁷ Recent evidence has shown

R. Mendoza et al.

beneficial effects of PNE in chronic musculoskeletal pain⁶⁹ and fibromyalgia.¹¹⁷

Exercise

Exercise can provide both acute and chronic changes in pain and BP.^{118–120} While exercise can increase BP during activity, there is a drop in BP for up to several hours after exercise in those with and without HTN.^{118,121} As suggested above, vitals should be measured throughout the plan of care to ensure safe delivery of exercise. Chronic adaptations in BP response to various forms of exercise are outlined in Supplementary Material 3. It should be noted that current data do not differentiate chronic changes in exercise between those with and without chronic pain. These changes should be used and interpreted with caution in patients with chronic pain.

Several mechanisms have been suggested as to why exercise is effective for managing chronic pain and HTN. Namely, there are improvements in pain regulation systems through EIH; improvements in autonomic regulation through decreasing sympathetic tone and vagal inhibition; increases in circulatory and intraarticular anti-inflammatory markers, and changes in brain chemical makeup, activation, and connectivity.¹²²

If deemed safe to begin an exercise program, factors such as current activity level, patient's previous response to exercise, patient beliefs, readiness to begin an exercise program, predominant pain mechanism (s), health conditions, and current medications need to be considered. Integrating exercise in the plan of care in those with chronic pain can be challenging. Nijs et al outline a stepwise approach to assist with implementing exercise for patients with chronic pain.¹²³ This includes an individually tailored plan that may involve:

- 1) Goal setting plus identifying body functions, activity limitations, participation restrictions, and contextual factors related to pain
- 2) Shared decision making
- 3) Tackling maladaptive beliefs about pain and physical activity
- Improving lifestyle factors that can optimize exercise, such as sleep, diet, and stress
- 5) Identification of activities that are avoided (managed with graded activity) versus persisted (managed with pacing and acceptancebased interventions).

Table 2 provides proposed guidelines for clinical-decision making regarding exercise parameters in patients with chronic pain and HTN.^{119,122,124–126} Aerobic, resistance, and isometric exercise have demonstrated effectiveness in managing patients with chronic pain or HTN.^{119,122,124–126} Exercise can be tailored to best meet the demands of the patient. For instance, if a patient has significant pain at the knee yet can tolerate regional exercise at other body sites, clinicians can consider titrating exercise at these other and less painful sites.¹²⁷

Moderate intensity aerobic exercise has been a staple element of HTN management at all stages.^{136,137} High intensity aerobic exercise is safe and also appears to promote substantial decreases in BP in those with HTN, although there are limited high quality studies supporting this approach.¹²⁸ Moreover, higher intensity training may not be tolerated in those with chronic pain, particularly early on during the rehabilitation process. It is important to recognize that, although exercise and lifestyle change are important management strategies in those with HTN, pharmacological intervention may be indicated in those with higher BP and/or those at higher risk for a cardiovascular event.⁹³ Moderate to high intensity aerobic exercise has also been recommended for the best analgesic effect. However, Vaegter and Jones²¹ suggested that even very short-duration aerobic exercise can elicit EIH, which implies that intensity, or permutations of intensity and duration, may be more critical for achieving analgesia after aerobic exercise than either variable alone.

Depending on the patient's pain and BP response to aerobic exercise/ testing, resistance and/or isometric exercise may be a necessary and

Table 2

Beginning clinical decision-making exercise parameter guidelines for those deemed safe to exercise with chronic pain^{113,128–131}.

Factor	Low-moderate intensity exercise	Moderate-high intensity exercise
Current activity level Previous exercise	Sedentary-low Negative	Moderate-high Positive
response Exercise/movement beliefs	Negative	Positive
Pain intensity/irritability Vitals at baseline	Moderate-high 130-179/80-109 mmHg for asymptomatic cases of HTN	Low <130/80 mm Hg
Vitals during submaximal aerobic activity	Vitals responding inappropriately to testing per metabolic equivalent	Vitals responding appropriately to testing per metabolic equivalent
Vitals post-aerobic activity	Elevated after five minutes of rest	Returning to baseline after five minutes of rest
Neuromusculoskeletal measures (range of motion)	High pain intensity High pain irritability Multiple motions reproduce pain	Low pain intensity Low pain irritability
Palpation Pain pressure thresholds Conditioned pain modulation	Hyperalgesia/allodynia Low No change or decreases	Normal response High Increases
Temporal summation Starting exercise parameters	High Aerobic ^	Low Aerobic
IntensitySets x repetitions.Frequency; Duration	 40-70 % HR max or 8- 13 rating of perceived exertion (RPE)- (RPE scale required if on beta blocker) 2-5x/week 20-60 minutes Resistance* 	 55-90 % HR max or 11-16 rating of perceived exertion (RPE scale required if on beta blocker) 4-7x/week 30-60 minutes Resistance
	 40-70 % of maximum 1-2 × 15-20 2-3x/week Isometric*^{132,133} 	 60-80 % of maximum 1-3 × 8-12 2x/week Isometric*¹³⁴
	 Up to 20 % of maximum 2-5s for 10 minutes 2-5x/week 	 Up to 80 % of maximum Up to 5 × 45s Up to 4x/week

[^]Consider shorter intervals to begin that add up to 20 minutes. ^{*} Consider starting with exercise at non-painful/least painful sites.^{119,135}

effective first step to address chronic pain and HTN. There is promising research emerging on the effects of isometric exercise and BP management.^{136,138} Large increases in BP can take place with resistance and isometric exercise.¹²¹ If there is a concern for large increases in BP, clinicians should consider using smaller musculature to begin.¹³⁸

Breathing/meditation

Emerging data suggest that meditation and/or breathing control exercises for 15-20 min/day can reduce BP in those with HTN ³⁹ and improve ability in those with chronic musculoskeletal pain.^{136,139} Research on mindfulness meditation has demonstrated significant pain relief and greater activation in brain regions associated with cognitive modulation of pain, including the orbitofrontal, subgenual anterior cingulate, and anterior insular cortex.¹⁴⁰

Manual therapy

Manual therapy intervention has shown to increase PPT and

facilitates impaired conditioned pain modulation in persons with lateral epicondylalgia of at least 6 weeks duration; however, the effects on BP were not reported.^{141,142} Yung et al found that non-thrust mobilization targeting the cervical spine in those with non-chronic neck pain results in acute changes in systolic BP.¹⁴³ Other trials are underway investigating the effects of manual therapy on BP in those with HTN.¹⁴⁴ While studies are limited, it seems plausible that improving pain may alter intertwined mechanisms involving chronic pain and HTN and reduce BP. Future studies are warranted.

Future directions

We advocate for more research to identify and understand pathophysiological mechanisms and factors underlying the relationship between chronic pain and HTN. Understanding these mechanisms should lead to more targeted and succinct evidence-informed management for patients with chronic pain and HTN. Future research should also ascertain patients whose HTN is directly linked to chronic pain, and if targeting chronic pain can result in improved BP regulation. Finally, we recommend best practice framework and guideline development to assist clinicians in managing the patient with chronic pain and HTN.

Conclusion

The personal and societal burden of chronic pain and HTN are significant. Although evidence suggests a relationship between chronic pain and HTN exists, there is much we have yet to learn. In conjunction with other health care providers, physical therapists can play a critical role in the management of patients with chronic pain and HTN. Identifying common risk factors, performing a thorough examination, and ongoing reassessment should allow progression of appropriate management that leads to successful outcomes for the patient with chronic pain and HTN.

Declaration of competing interest

The author has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.bjpt.2025.101229.

References

- 1. Hypertension. Accessed August 29th, 2023, https://www.who.int/news-room/fact-sheets/detail/hypertension.
- Giummarra MJ, Tardif H, Blanchard M, Tonkin A, Arnold CA. Hypertension prevalence in patients attending tertiary pain management services, a registrybased Australian cohort study. *PLoS One*. 2020;15(1), e0228173.
- Zis P, Daskalaki A, Bountouni I, Sykioti P, Varrassi G, Paladini A. Depression and chronic pain in the elderly: links and management challenges. *Clin Interv Aging*. 2017:709–720.
- Khalid S, Sambamoorthi U, Umer A, Lilly CL, Gross DK, Innes KE. Increased odds of incident Alzheimer's disease and related dementias in presence of common noncancer chronic pain conditions in Appalachian older adults. J Aging Health. 2022; 34(2):158–172.
- Gaskin DJ, Richard P. The economic costs of pain in the United States. J Pain. 2012;13(8):715–724.
- Ditre JW, Zale EL, LaRowe LR. A reciprocal model of pain and substance use: Transdiagnostic considerations, clinical implications, and future directions. *Annu Rev Clin Psychol.* 2019;15(1):503–528.
- NIo Health. Interagency Pain Research Coordinating Committee: National Pain Strategy: A comprehensive population health-level strategy for pain. US Department of Health and Human Services, Published; 2016:2023.
- BoHS Policy, Research CoAP Care. Relieving pain in America: a blueprint for transforming prevention, care, education, and research. National Academies Press; 2011.

- Kirkland EB, Heincelman M, Bishu KG, et al. Trends in healthcare expenditures among US adults with hypertension: national estimates, 2003–2014. J Am Heart Assoc. 2018;7(11), e008731.
- Shrestha S, Bauer CX, Hendricks B, Stopka TJ. Spatial epidemiology: an empirical framework for syndemics research. Soc Sci Med (1967). 2022;295, 113352.
- Chung OY, Bruehl S, Diedrich L, Diedrich A, Chont M, Robertson D. Baroreflex sensitivity associated hypoalgesia in healthy states is altered by chronic pain. *Pain*. 2008;138(1):87–97.
- Goodson NJ, Smith BH, Hocking LJ, et al. Cardiovascular risk factors associated with the metabolic syndrome are more prevalent in people reporting chronic pain: results from a cross-sectional general population study. *PAIN*®. 2013;154(9): 1595–1602.
- LaRowe LR, Ditre JW. Pain, nicotine, and tobacco smoking: Current state of the science. Pain. 2020;161(8):1688.
- Li C-Y, Lin W-C, Lu C-Y, Chung YS, Cheng Y-C. Prevalence of pain in communitydwelling older adults with hypertension in the United States. *Sci Rep.* 2022;12(1): 1–10.
- Heikkala E, Oura P, Ho E, Ferreira P, Paananen M, Karppinen J. Accumulation of long-term diseases is associated with musculoskeletal pain dimensions among middle-aged individuals with musculoskeletal pain. *Eur J Pain*. 2023;27(3): 438-448.
- 16. Ramanathan S, Hibbert P, Wiles L, Maher CG, Runciman W. What is the association between the presence of comorbidities and the appropriateness of care for low back pain? A population-based medical record review study. *BMC Musculoskelet Disord*. 2018;19:1–9.
- Warwick H, O'Donnell J, Mather III RC, Jiranek W. Disparity of health services in patients with knee osteoarthritis before total knee arthroplasty. *Arthroplast Today*. 2020;6(1):81–87.
- Bruehl S, Olsen RB, Tronstad C, et al. Chronic pain-related changes in cardiovascular regulation and impact on comorbid hypertension in a general population: the Tromsø study. *Pain*. 2018;159(1):119–127.
- Olsen RB, Bruehl S, Nielsen CS, Rosseland LA, Eggen AE, Stubhaug A. Hypertension prevalence and diminished blood pressure-related hypoalgesia in individuals reporting chronic pain in a general population: the Tromsø study. *Pain.* Feb 2013; 154(2):257–262. https://doi.org/10.1016/j.pain.2012.10.020.
- Levy M, Pappano A, Berne R. Cardiovascular physiology. 9. Philadelphia, PA: Mosby. Elsevier; 2007.
- Vaegter HB, Jones MD. Exercise-induced hypoalgesia after acute and regular exercise: experimental and clinical manifestations and possible mechanisms in individuals with and without pain. *Pain Rep.* 2020;5(5).
- Ring C, Edwards L, Kavussanu M. Effects of isometric exercise on pain are mediated by blood pressure. *Biol Psychol.* 2008;78(1):123–128.
- Chimenti RL, Frey-Law LA, Sluka KA. A mechanism-based approach to physical therapist management of pain. *Phys Ther.* 2018;98(5):302–314. https://doi.org/ 10.1093/pti/pzv030.
- 24. Pereira-Silva R, Costa-Pereira JT, Alonso R, Serrão P, Martins I, Neto FL. Attenuation of the diffuse noxious inhibitory controls in chronic joint inflammatory pain is accompanied by anxiodepressive-like behaviors and impairment of the descending noradrenergic modulation. *Int J Mol Sci.* 2020;21(8): 2973.
- Hayashida K-i, Obata H. Strategies to treat chronic pain and strengthen impaired descending noradrenergic inhibitory system. Int J Mol Sci. 2019;20(4):822.
- Bruehl S, Chung OY. Interactions between the cardiovascular and pain regulatory systems: an updated review of mechanisms and possible alterations in chronic pain. *Neurosci Biobehav Rev.* 2004;28(4):395–414.
- Hering L, Rahman M, Potthoff SA, Rump LC, Stegbauer J. Role of α2-adrenoceptors in hypertension: focus on renal sympathetic neurotransmitter release, inflammation, and sodium homeostasis. *Front Physiol.* 2020;11, 566871.
- Hsu C-N, Tain Y-L. Targeting the renin–angiotensin–aldosterone system to prevent hypertension and kidney disease of developmental origins. *Int J Mol Sci.* 2021;22 (5):2298.
- 29. Capric V, Chandrakumar HP, Celenza-Salvatore J, Makaryus AN. The role of the renin-angiotensin-aldosterone system in cardiovascular disease: pathogenetic insights and clinical implications. *Renin-Angiotensin Aldosterone Syst.* 2021.
- 30. Pertovaara A. Noradrenergic pain modulation. Prog Neurobiol. 2006;80(2):53-83.
- Drummond ES, Dawson LP, Finch PM, Bennett GJ, Drummond PD. Increased expression of cutaneous α1-adrenoceptors after chronic constriction injury in rats. *J Pain*. 2014;15(2):188–196.
- Fuchs PN, Meyer RA, Raja SN. Heat, but not mechanical hyperalgesia, following adrenergic injections in normal human skin. *Pain.* 2001;90(1-2):15–23.
- Dogrul A, Coskun I, Uzbay T. The contribution of alpha-1 and alpha-2 adrenoceptors in peripheral imidazoline and adrenoceptor agonist-induced nociception. *Anesth Analg.* 2006;103(2):471–477.
- Carroll I, Mackey S, Gaeta R. The role of adrenergic receptors and pain: The good, the bad, and the unknown. Elsevier; 2007:17–21.
- 35. Gil DW, Cheevers CV, Kedzie KM, et al. α-1-Adrenergic receptor agonist activity of clinical α-adrenergic receptor agonists interferes with α-2-mediated analgesia. *J Am Soc Anesthesiol*. 2009;110(2):401–407.
- Suarez-Roca H, Klinger RY, Podgoreanu MV, et al. Contribution of baroreceptor function to pain perception and perioperative outcomes. *Anesthesiology*. 2019;130 (4):634–650.
- Kaufmann H, Norcliffe-Kaufmann L, Dysfunction Palma J-ABaroreflex. N Engl J Med. 2020;382(2):163–178. https://doi.org/10.1056/NEJMra1509723.
- Virtanen R, Jula A, Salminen JK, et al. Anxiety and hostility are associated with reduced baroreflex sensitivity and increased beat-to-beat blood pressure

R. Mendoza et al.

variability. Psychosom Med. 2003;65(5):751–756. https://doi.org/10.1097/01. Psy.0000088760.65046.Cf.

- Okifuji A, Bradshaw DH, Olson C. Evaluating obesity in fibromyalgia: neuroendocrine biomarkers, symptoms, and functions. *Clin Rheumatol.* 2009;28: 475–478.
- Briggs MS, Givens DL, Schmitt LC, Taylor CA. Relations of C-reactive protein and obesity to the prevalence and the odds of reporting low back pain. *Arch Phys Med Rehabil.* 2013;94(4):745–752.
- Zewinger S, Reiser J, Jankowski V, et al. Apolipoprotein C3 induces inflammation and organ damage by alternative inflammasome activation. *Nat Immunol.* 2020;21 (1):30–41.
- **42.** Guzik TJ, Touyz RM. Oxidative stress, inflammation, and vascular aging in hypertension. *Hypertension*. 2017;70(4):660–667.
- Anderson L, Nguyen TT, Dall CH, Burgess L, Bridges C, Taylor RS. Exercise-based cardiac rehabilitation in heart transplant recipients. *Cochrane Database Syst Rev.* Apr 4 2017;4, CD012264. https://doi.org/10.1002/14651858.CD012264.pub2.
- Fang X-X, Zhai M-N, Zhu M, et al. Inflammation in pathogenesis of chronic pain: foe and friend. *Mol Pain*. 2023;19, 17448069231178176.
- Aboukhater D, Morad B, Nasrallah N, et al. Inflammation and hypertension: underlying mechanisms and emerging understandings. J Cell Physiol. 2023;238(6): 1148–1159.
- Baral P, Udit S, Chiu IM. Pain and immunity: implications for host defence. Nat Rev Immunol. 2019;19(7):433–447.
- Zhuo M. Neural mechanisms underlying anxiety-chronic pain interactions. *Trends Neurosci.* 2016;39(3):136–145.
- 48. AACVPR. Guidelines for cardiac rehabilitation programs. Human Kinetics. 2020.
- Delitto A, George SZ, Van Dillen L, et al. Low back pain: clinical practice guidelines linked to the international classification of functioning, disability, and health from the orthopaedic section of the American physical therapy association. *J Orthop Sports Phys Ther.* 2012;42(4):A1–A57. https://doi.org/10.2519/jospt.2012.42.4. A1.
- Tait RC, Chibnall JT, House K, Biehl J. Medical judgments across the range of reported pain severity: clinician and lay perspectives. *Pain Med.* 2016;17(7): 1269–1281.
- De Ruddere L, Goubert L, Stevens M, ACdC Williams, Crombez G. Discounting pain in the absence of medical evidence is explained by negative evaluation of the patient. *PAIN*[®]. 2013;154(5):669–676.
- Hensley CP, Emerson AJ. Non-small cell lung carcinoma: clinical reasoning in the management of a patient referred to physical therapy for costochondritis. *Phys Ther.* 2018;98(6):503–509.
- Carruthers B, van de Sande M, De Meirleir K, et al. Myalgic encephalomyelitis: international consensus criteria (vol 270, pg 327, 2011). J Intern Med. 2017;282 (4), 353-353.
- Jason LA, Dorri JA. ME/CFS and post-exertional malaise among patients with long COVID. Neurol Int. 2022;15(1):1–11.
- Vernon SD, Hartle M, Sullivan K, et al. Post-exertional malaise among people with long COVID compared to myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). Work. 2023;74(4):1179–1186.
- Davis HE, Assaf GS, McCorkell L, et al. Characterizing long COVID in an international cohort: 7 months of symptoms and their impact. *EClinicalMedicine*. 2021;38.
- Twomey R, DeMars J, Franklin K, SN Culos-Reed, Weatherald J, Wrightson JG. Chronic fatigue and postexertional malaise in people living with long COVID: an observational study. *Phys Ther.* 2022;102(4):pzac005.
- Bai NA, Richardson CS. Posttreatment Lyme disease syndrome and myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review and comparison of pathogenesis. *Chronic Dis Transl Med.* 2023;9(03):183–190.
- Moore GE, Keller BA, Stevens J, et al. Recovery from exercise in persons with myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *Medicina (B Aires)*. 2023;59(3):571.
- 60. Keller B, Receno CN, Franconi CJ, et al. Cardiopulmonary and metabolic responses during a 2-day CPET in myalgic encephalomyelitis/chronic fatigue syndrome: translating reduced oxygen consumption to impairment status to treatment considerations. J Transl Med. 2024;22(1):1–35.
- Mateo LJ, Chu L, Stevens S, et al. Post-exertional symptoms distinguish myalgic encephalomyelitis/chronic fatigue syndrome subjects from healthy controls. Work. 2020;66(2):265–275.
- 62. Franklin JD, Graham M. Repeated maximal exercise tests of peak oxygen consumption in people with myalgic encephalomyelitis/chronic fatigue syndrome: a systematic review and meta-analysis. *Fatigue: Biomed Health Behav.* 2022;10(3): 119–135.
- Lim E-J, Kang E-B, Jang E-S, Son C-G. The prospects of the two-day cardiopulmonary exercise test (CPET) in ME/CFS patients: a meta-analysis. J Clin Med. 2020;9(12):4040.
- Davenport TE, Stevens SR, VanNess MJ, Snell CR, Little T. Conceptual model for physical therapist management of chronic fatigue syndrome/myalgic encephalomyelitis. *Phys Ther.* 2010;90(4):602–614.
- 65. Shraim MA, Sluka KA, Sterling M, et al. Features and methods to discriminate between mechanism-based categories of pain experienced in the musculoskeletal system: a Delphi expert consensus study. *Pain*. 2022;163(9):1812–1828.
- 66. Nijs J, Kosek E, Chiarotto A, et al. Nociceptive, neuropathic, or nociplastic low back pain? the low back pain phenotyping (BACPAP) consortium's international and multidisciplinary consensus recommendations. *Lancet Rheumatol.* Mar 2024;6 (3):e178–e188. https://doi.org/10.1016/s2665-9913(23)00324-7.
- 67. Hensley CP, Courtney CA. Management of a patient with chronic low back pain and multiple health conditions using a pain mechanisms-based classification

Brazilian Journal of Physical Therapy 29 (2025) 101229

approach. J Orthop Sports Phys Ther. Jun 2014;44(6). https://doi.org/10.2519/ jospt.2014.4861, 403-c2.

- Nijs J, Van Wilgen CP, Van Oosterwijck J, van Ittersum M, Meeus M. How to explain central sensitization to patients with 'unexplained'chronic musculoskeletal pain: practice guidelines. *Man Ther.* 2011;16(5):413–418.
- 69. Siddal B, Ram A, Jones MD, Booth J, Perriman D, Summers SJ. Short-term impact of combining pain neuroscience education with exercise for chronic musculoskeletal pain: a systematic review and meta-analysis. *Pain*. 2022;163(1): e20–e30.
- Neblett R, Cohen H, Choi Y, et al. The Central Sensitization Inventory (CSI): establishing clinically significant values for identifying central sensitivity syndromes in an outpatient chronic pain sample. J Pain. 2013;14(5):438–445.
- Freynhagen R, Tölle TR, Gockel U, Baron R. The painDETECT project-far more than a screening tool on neuropathic pain. *Curr Med Res Opin.* 2016;32(6): 1033–1057.
- 72. Freynhagen R, Baron R, Gockel U, Tölle TR. Pain DETECT: a new screening questionnaire to identify neuropathic components in patients with back pain. *Curr Med Res Opin*. 2006;22(10):1911–1920.
- Mathieson S, Maher CG, Terwee CB, De Campos TF, Lin C-WC. Neuropathic pain screening questionnaires have limited measurement properties. A systematic review. J Clin Epidemiol. 2015;68(8):957–966.
- 74. Bennett M. Leeds Assessment of neuropathic symptoms and signs pain scale. Pain.
- Backonja MM, Attal N, Baron R, et al. Value of quantitative sensory testing in neurological and pain disorders: NeuPSIG consensus. *Pain.* Sep 2013;154(9): 1807–1819. https://doi.org/10.1016/j.pain.2013.05.047.
- 76. Huysmans E, Goudman L, Van Bogaert W, et al. Experimental pain measurements do not relate to pain intensity and pain cognitions in people scheduled for surgery for lumbar radiculopathy. *Pain Med.* 2023;24(2):139–149.
- Palsson TS, Christensen W, De Martino E, Graven-Nielsen T. Pain and disability in low back pain can be reduced despite no significant improvements in mechanistic pain biomarkers. *Clin J Pain*. 2021;37(5):330–338.
- Balasch-Bernat M, Lluch E, Vaegter H, Dueñas L. Should exercises be painful or not? Effects on clinical and experimental pain in individuals with shoulder pain. *J Pain*. 2021;22(10):1246–1255.
- 79. Sachau J, Appel C, Reimer M, et al. Test–retest reliability of a simple bedsidequantitative sensory testing battery for chronic neuropathic pain. *Pain Rep.* 2023;8 (1):e1049.
- Zhu GC, Böttger K, Slater H, et al. Concurrent validity of a low-cost and timeefficient clinical sensory test battery to evaluate somatosensory dysfunction. *Eur J Pain*. 2019;23(10):1826–1838.
- Reimer M, Forstenpointner J, Hartmann A, et al. Sensory bedside testing: a simple stratification approach for sensory phenotyping. *Pain Rep.* May-Jun 2020;5(3): e820. https://doi.org/10.1097/pr9.00000000000820.
- Coppieters I, Meeus M, Kregel J, et al. Relations Between Brain Alterations and Clinical Pain Measures in Chronic Musculoskeletal Pain: A Systematic Review. J Pain. Sep 2016;17(9):949–962. https://doi.org/10.1016/j.jpain.2016.04.005.
- O'Leary H, Smart KM, Moloney NA, Blake C, Doody CM. Pain sensitization associated with nonresponse after physiotherapy in people with knee osteoarthritis. *Pain*. Sep 2018;159(9):1877–1886. https://doi.org/10.1097/j. pain.000000000001288.
- 84. Vaegter HB, Petersen KK, Mørch CD, Imai Y, Arendt-Nielsen L. Assessment of CPM reliability: quantification of the within-subject reliability of 10 different protocols. *Scand J Pain*. Oct 25 2018;18(4):729–737. https://doi.org/10.1515/sjpain-2018-0087.
- Wideman TH, Edwards RR, Finan PH, Haythornthwaite JA, Smith MT. Comparing the predictive value of task performance and task-specific sensitivity during physical function testing among people with knee osteoarthritis. *J Orthop Sports Phys Ther.* May 2016;46(5):346–356. https://doi.org/10.2519/jospt.2016.6311.
- Muntner P, Shimbo D, Carey RM, et al. Measurement of blood pressure in humans: a scientific statement from the American heart association. *Hypertension*. 2019;73 (5):e35–e66.
- **87.** Cifu AS, Davis AM. Prevention, detection, evaluation, and management of high blood pressure in adults. *JAMA*. 2017;318(21):2132–2134.
- 88. Sethi R, Hiremath J, Ganesh V, et al. Correlation between stroke risk and systolic blood pressure in patients over 50 years with uncontrolled hypertension: results from the SYSTUP-India study. *Cardiovasc Ther.* 2021;2021(1), 6622651.
- 89. Miyamoto GC, Ben AJ, Bosmans JE, et al. Interpretation of trial-based economic evaluations of musculoskeletal physical therapy interventions. *Braz J Phys Ther*. 2021;25(5):514–529.
- 90. Control CoD. About Stroke. center of disease control https://www.cdc.gov/stro ke/about/index.html.
- 91. Whelton PK, Carey RM, Aronow WS, et al. ACC/AHA/AAPA/ABC/ACPM/AGS/ APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. J Am Coll Cardiol. 2017;71(19):e127–e248, 2018.
- **92.** Adler J, Dekerlegand J. Implementation of vital sign and activity guidelines for the outpatient therapist. *Patient Safety*. 2023;5(3), 88307.
- Unger T, Borghi C, Charchar F, et al. 2020 International Society of Hypertension global hypertension practice guidelines. *Hypertension*. 2020;75(6):1334–1357.
- **94.** Paluch AE, Boyer WR, Franklin BA, et al. Resistance exercise training in individuals with and without cardiovascular disease: 2023 update: a scientific statement from the American Heart Association. *Circulation*. 2024;149(3):e217–e231.
- Ghadieh AS, Saab B. Evidence for exercise training in the management of hypertension in adults. *Can Fam Phys.* Mar 2015;61(3):233–239.

R. Mendoza et al.

- **96.** Janke AT, McNaughton CD, Brody AM, Welch RD, Levy PD. Trends in the incidence of hypertensive emergencies in US emergency departments from 2006 to 2013. *J Am Heart Assoc.* 2016;5(12), e004511.
- Siddiqi TJ, Rashid AM, Siddiqi AK, et al. Association of electronic cigarette exposure on cardiovascular health: a systematic review and meta-analysis. *Curr Probl Cardiol.* 2023;48(9), 101748.
- 98. Schultz MG, Otahal P, Cleland VJ, Blizzard L, Marwick TH, Sharman JE. Exerciseinduced hypertension, cardiovascular events, and mortality in patients undergoing exercise stress testing: a systematic review and meta-analysis. *Am J Hypertens*. 2013;26(3):357–366.
- Weiss SA, Blumenthal RS, Sharrett AR, Redberg RF, Mora S. Exercise blood pressure and future cardiovascular death in asymptomatic individuals. *Circulation*. May 18 2010;121(19):2109–2116. https://doi.org/10.1161/ circulationaba.109.895292.
- 100. Mariampillai JE, Liestøl K, Kjeldsen SE, et al. Exercise systolic blood pressure at moderate workload is linearly associated with coronary disease risk in healthy men. *Hypertension*. 2020;75(1):44–50.
- 101. Zafrir B, Aker A, Asaf Y, Saliba W. Blood pressure response during treadmill exercise testing and the risk for future cardiovascular events and new-onset hypertension. J Hypertens. 2022;40(1):143–152.
- 102. McHam SA, Marwick TH, Pashkow FJ, Lauer MS. Delayed systolic blood pressure recovery after graded exercise: an independent correlate of angiographic coronary disease. J Am Coll Cardiol. 1999;34(3):754–759.
- 103. Kokkinos P, Kaminsky LA, Arena R, Zhang J, Myers J. New generalized equation for predicting maximal oxygen uptake (from the Fitness Registry and the Importance of Exercise National Database). Am J Cardiol. 2017;120(4):688–692.
- 104. Eston R, Connolly D. The use of ratings of perceived exertion for exercise prescription in patients receiving β-blocker therapy. *Sports Med.* 1996;21:176–190.
- 105. Peterman JE, Arena R, Myers J, et al. Reference Standards for Cardiorespiratory Fitness by Cardiovascular Disease Category and Testing Modality: Data From FRIEND. J Am Heart Assoc. 2021;10(22), e022336. https://doi.org/10.1161/ JAHA.121.022336.
- 106. Fletcher GF, Ades PA, Kligfield P, et al. Exercise standards for testing and training: a scientific statement from the American Heart Association. *Circulation*. 2013;128 (8):873–934.
- 107. Pickering TG, Hall JE, Appel LJ, et al. Recommendations for blood pressure measurement in humans and experimental animals: Part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension*. Jan 2005;45(1):142–161. https://doi. org/10.1161/01.HYP.000150859.47929.8e.
- 108. Schultz MG, Currie KD, Hedman K, et al. The identification and management of high blood pressure using exercise blood pressure: current evidence and practical guidance. *Int J Environ Res Public Health*. 2022;19(5):2819.
- Minhas H, Morton C, Shaw M, Shelley B. Reproducibility of heart rate recovery measures across differing modalities and intensities of submaximal exercise in healthy volunteers: a healthy volunteer study. *Anesthesiol Perioperat Sci.* 2024;2(1): 11. https://doi.org/10.1007/s44254-023-00042-2, 2024/02/26.
- 110. Bruneau Jr ML, Johnson BT, Huedo-Medina TB, Larson KA, Ash GI, Pescatello LS. The blood pressure response to acute and chronic aerobic exercise: A meta-analysis of candidate gene association studies. J Sci Med Sport. 2016;19(5):424–431.
- 111. Laukkanen JA, Kurl S, Salonen R, Lakka TA, Rauramaa R, Salonen JT. Systolic blood pressure during recovery from exercise and the risk of acute myocardial infarction in middle-aged men. *Hypertension*. 2004;44(6):820–825.
- **112.** Wang JJ, Zheng Z, Zhang Y. Association of postexercise blood pressure with cardiovascular outcomes and mortality: The CARDIA study. *J Clin Hypertens*. 2024.
- 113. Cataldo D, Anthony Figueroa, Robinson B, Ryan Sandstrom. Exercising your way to lowering your blood pressure. Am Coll Sport Med https://www.acsm.org/docs/def ault-source/files-for-resource-library/exercising-your-way-to-lowering-your-bloo d-pressure.pdf?sfvrsn=1aa17ebd_2.
- Rivasi G, Menale S, Turrin G, Coscarelli A, Giordano A, Ungar A. The effects of pain and analgesic medications on blood pressure. *Curr Hypertens Rep.* 2022;24(10): 385–394.
- 115. Nijs J, Wyns A, Hendrix J. The importance of stress in the paradigm shift from a tissue-and disease-based pain management approach towards multimodal lifestyle interventions for chronic pain. *Braz J Phys Ther.* 2024;28(2).
- 116. Nijs J, Mairesse O, Tang NK. The importance of sleep in the paradigm shift from a tissue-and disease-based pain management approach towards multimodal lifestyle interventions for chronic pain. *Braz J Phys Ther.* 2024;28(1).
- 117. Van Oosterwijck J, Meeus M, Paul L, et al. Pain physiology education improves health status and endogenous pain inhibition in fibromyalgia: a double-blind randomized controlled trial. *Clin J Pain*. Oct 2013;29(10):873–882. https://doi. org/10.1097/AJP.0b013e31827c7a7d.
- Gkaliagkousi E, Gavriilaki E, Douma S. Effects of acute and chronic exercise in patients with essential hypertension: benefits and risks. *Am J Hypertens*. Apr 2015; 28(4):429–439. https://doi.org/10.1093/ajh/hpu203.
- 119. Rice D, Nijs J, Kosek E, et al. Exercise-induced hypoalgesia in pain-free and chronic pain populations: state of the art and future directions. *J Pain*. 2019;20(11): 1249–1266.

- Brazilian Journal of Physical Therapy 29 (2025) 101229
- Lopes S, Afreixo V, Teixeira M, et al. Exercise training reduces arterial stiffness in adults with hypertension: a systematic review and meta-analysis. J Hypertens. Feb 1 2021;39(2):214–222. https://doi.org/10.1097/hjh.00000000002619.
- 121. Edwards JJ, Deenmamode AH, Griffiths M, et al. Exercise training and resting blood pressure: a large-scale pairwise and network meta-analysis of randomised controlled trials. *Br J Sports Med.* 2023;57(20):1317–1326.
- 122. Sluka KA, Law LF, Bement MH. Exercise-induced pain and analgesia? Underlying mechanisms and clinical translation. *Pain*. 2018;159(Suppl 1):S91.
- 123. Nijs J, Mostaqim K, Oostendorp R. Exercise therapy and physical activity in the paradigm shift from a tissue-and disease-based pain management approach towards multimodal lifestyle interventions for chronic pain. *Braz J Phys Ther.* 2024; 28(4).
- 124. de Zoete RM, Chen K, Sterling M. Central neurobiological effects of physical exercise in individuals with chronic musculoskeletal pain: a systematic review. *BMJ Open.* 2020;10(7), e036151.
- 125. Yeater TD, Cruz CJ, Cruz-Almeida Y, Allen KD. Autonomic nervous system dysregulation and osteoarthritis pain: mechanisms, measurement, and future outlook. *Curr Rheumatol Rep.* 2022;24(6):175–183.
- 126. Puts S, Liberman K, Leysen L, et al. Exercise-induced effects on inflammatory markers and brain-derived neurotrophic factor in patients with knee osteoarthritis. A systematic review with meta-analysis. *Exerc Immunol Rev.* 2023;29:22–53.
- 127. Alsouhibani A, Hoeger Bement M. Impaired conditioned pain modulation was restored after a single exercise session in individuals with and without fibromyalgia. *Pain Rep.* May-Jun 2022;7(3):e996. https://doi.org/10.1097/ pr9.000000000000996.
- 128. Teixeira JM, Motta-Santos D, Milanovic Z, Pereira RL, Krustrup P, Povoas S. Intermittent high-intensity exercise for pre-to established hypertension: A systematic review and meta-analysis. *Scand J Med Sci Sports*, 2023;33(4):364–381.
- 129. Sharman JE, Smart NA, Coombes JS, Stowasser M. Exercise and sport science australia position stand update on exercise and hypertension. J Hum Hypertens. 2019;33(12):837–843.
- 130. Ferro Moura Franco K, Lenoir D, dos Santos Franco YR, Jandre Reis FJ, Nunes Cabral CM, Meeus M. Prescription of exercises for the treatment of chronic pain along the continuum of nociplastic pain: A systematic review with meta-analysis. *Eur J Pain*. 2021;25(1):51–70.
- 131. Correia RR, Veras ASC, Tebar WR, Rufino JC, Batista VRG, Teixeira GR. Strength training for arterial hypertension treatment: a systematic review and meta-analysis of randomized clinical trials. *Sci Rep.* 2023;13(1):201.
- Vuvan V, Vicenzino B, Mellor R, Heales LJ, Coombes BK. Unsupervised isometric exercise versus wait-and-see for lateral elbow tendinopathy. *Med Sci Sports Exerc*. 2020;52(2):287–295.
- **133.** Berardi G, Senefeld JW, Hunter SK, Bernent MKH. Impact of isometric and concentric resistance exercise on pain and fatigue in fibromyalgia. *Eur J Appl Physiol.* 2021;121:1389–1404.
- 134. Lim HY, Wong SH. Effects of isometric, eccentric, or heavy slow resistance exercises on pain and function in individuals with patellar tendinopathy: a systematic review. *Physiother. Res. Int.* 2018;23(4):e1721.
- 135. Christensen N, Black L, Furze J, Huhn K, Vendrely A, Wainwright S. Clinical reasoning: survey of teaching methods, integration, and assessment in entry-level physical therapist academic education. *Phys Ther.* 2017;97(2):175–186.
- 136. Fu J, Liu Y, Zhang L, et al. Nonpharmacologic interventions for reducing blood pressure in adults with prehypertension to established hypertension. J Am Heart Assoc. 2020;9(19), e016804.
- 137. Hanssen H, Boardman H, Deiseroth A, et al. Personalized exercise prescription in the prevention and treatment of arterial hypertension: a Consensus Document from the European Association of Preventive Cardiology (EAPC) and the ESC Council on Hypertension. *Eur J Prev Cardiol.* 2022;29(1):205–215.
- **138.** Hansford HJ, Parmenter BJ, McLeod KA, et al. The effectiveness and safety of isometric resistance training for adults with high blood pressure: a systematic review and meta-analysis. *Hypertens Res.* 2021;44(11):1373–1384.
- 139. Lin T-H, Tam K-W, Yang Y-L, Liou T-H, Hsu T-H, Rau C-L. Meditation-based therapy for chronic low back pain management: a systematic review and metaanalysis of randomized controlled trials. *Pain Med.* 2022;23(10):1800–1811.
- 140. Zeidan F, Emerson NM, Farris SR, et al. Mindfulness meditation-based pain relief employs different neural mechanisms than placebo and sham mindfulness meditation-induced analgesia. *J Neurosci.* 2015;35(46):15307–15325.
- Voogt L, de Vries J, Meeus M, Struyf F, Meuffels D, Nijs J. Analgesic effects of manual therapy in patients with musculoskeletal pain: a systematic review. *Man Ther.* Apr 2015;20(2):250–256. https://doi.org/10.1016/j.math.2014.09.001.
- **142.** Muhsen A, Moss P, Gibson W, et al. The association between conditioned pain modulation and manipulation-induced analgesia in people with lateral epicondylalgia. *Clin J Pain*. 2019;35(5):435–442.
- 143. Yung E, Oh C, Wong M, et al. Non-thrust cervical manipulations reduce short-term pain and decrease systolic blood pressure during intervention in mechanical neck pain: a randomized clinical trial. J Manual Manipulat Ther. 2020;28(2):82–93.
- Pederson E. The Immediate Effect of an Osteopathic Manual Treatment on Blood Pressure in Adults With Hypertension - a Pilot Study. https://clinicaltrials.gov/s tudy/NCT04479384.